

FATTENING AND NONFATTENING PHENOMENA FOR PLANAR NONLOCAL CURVATURE FLOWS

ANNALISA CESARONI, SERENA DIPIERRO, MATTEO NOVAGA, AND ENRICO VALDINOCI

ABSTRACT. We discuss fattening phenomenon for the evolution of planar curves according to their nonlocal curvature. More precisely, we consider a class of generalized curvatures which correspond to the first variation of suitable nonlocal perimeter functionals, defined in terms of an interaction kernel K , which is symmetric, nonnegative, possibly singular at the origin, and satisfies appropriate integrability conditions.

We prove a general result about uniqueness of the geometric evolutions starting from regular sets with positive K -curvature and we discuss the fattening phenomenon for the evolution starting from the cross, showing that this phenomenon is very sensitive to the strength of the interactions. As a matter of fact, we show that the fattening of the cross occurs for kernels with sufficiently large mass near the origin, while for kernels that are sufficiently weak near the origin such a fattening phenomenon does not occur.

We also provide some further results in the case of the fractional mean curvature flow, showing that strictly starshaped sets have a unique geometric evolution.

Moreover, we exhibit two illustrative examples of closed nonregular curves, the first with a Lipschitz-type singularity and the second with a cusp-type singularity, given by two tangent circles of equal radius, whose evolution develops fattening in the first case, and is uniquely defined in the second, thus remarking the high sensitivity of the fattening phenomenon in terms of the regularity of the initial datum. The latter example is in striking contrast to the classical case of the (local) curvature flow, where two tangent circles always develop fattening.

As a byproduct of our analysis, we provide also a simple proof of the fact that the cross is not a K -minimal set for the nonlocal perimeter functional associated to K .

CONTENTS

1. Introduction	2
1.1. Evolution of the cross	4
1.2. A remark on K -minimal cones	6
1.3. Fractional curvature evolution of starshaped sets	6
2. Regular sets of positive K -curvature and proof of Theorem 1.2	9
3. K -curvature of the perturbed cross and proofs of Proposition 1.4 and of Theorem 1.5	10
4. Moving boxes, weak interaction kernels and proof of Theorem 1.7	14
5. K -minimal cones and proof of Proposition 1.9	18
6. Strictly starshaped domains and proof of Proposition 1.10	20
7. Perturbed double droplet and proof of Theorem 1.11	20
8. Perturbation of tangent balls and proof of Theorem 1.13	23
Appendix A. Viscosity solutions and geometric barriers	25
References	32

2010 *Mathematics Subject Classification.* 53C44, 35D40, 35R11 58E12 .

Key words and phrases. Nonlocal curvature flows, planar curves, uniqueness of the geometric evolution, fattening phenomenon, nonlocal perimeter functionals, minimal sets.

Annalisa Cesaroni: Dipartimento di Scienze Statistiche, Università di Padova, Via Battisti 241/243, 35121 Padova, Italy. annalisa.cesaroni@unipd.it

Serena Dipierro: Department of Mathematics and Statistics, University of Western Australia, 35 Stirling Hwy, Crawley WA 6009, Australia, Dipartimento di Matematica, Università di Milano, Via Saldini 50, 20133 Milan, Italy. serydipierro@yahoo.it

Matteo Novaga: Dipartimento di Matematica, Università di Pisa, Largo Pontecorvo 5, 56127 Pisa, Italy. matteo.novaga@unipi.it

Enrico Valdinoci: Department of Mathematics and Statistics, University of Western Australia, 35 Stirling Hwy, Crawley WA 6009, Australia, School of Mathematics and Statistics, University of Melbourne, 813 Swanston St, Parkville VIC 3010, Australia, Dipartimento di Matematica, Università di Milano, Via Saldini 50, 20133 Milan, Italy. enrico@math.utexas.edu

1. INTRODUCTION

In this paper we are interested in the analysis of the fattening phenomenon for evolutions of sets in the plane according to nonlocal curvature flows. Fattening is a particular kind of singularity which arises in the evolution of boundaries by their (local or nonlocal) curvatures and more generally in geometric evolution of manifolds and is related to nonuniqueness of geometric solutions to the flow. Fattening phenomenon has been studied for mean curvature flow since long time and a complete characterization of initial data which develop fattening is still missing. In the case of the plane, it is known that smooth compact level curves never develop an interior, due to a result by Grayson on the evolution of regular compact curves. This result is no more valid for fractional mean curvature flow in the plane, as proved recently in [12]. We recall that examples of fattening of nonregular or noncompact curves in the plane for the mean curvature flow have been given in [3, 13, 15], where, in particular, the fattening of the evolution starting from the cross is proved. Finally nonfattening for strictly starshaped initial data is proved in [20], whereas nonfattening of convex and mean convex initial data is proved in [1], see also [2] and [4].

In this paper we start the analysis of the fattening phenomenon in the plane for general nonlocal curvature flows. This problem has not yet been considered in the literature apart from the result in [9] about nonfattening for convex initial data under fractional mean curvature evolution in any space dimension.

Here we will show that some results which are true for the mean curvature flow are still valid, such as nonfattening for regular initial data with positive curvature or strictly starshaped initial data.

Nevertheless, in general, some different behaviors with respect to the mean curvature flow arise, due to the fact that the fattening phenomenon is very sensitive to the strength of the nonlocal interactions. We discuss in particular the evolution starting from the cross, which develops fattening only if the interactions are sufficiently strong. Moreover, we show an example of a bounded set with positive curvature which fattens, and an example of a bounded set whose evolution by fractional mean curvature flow does not present fattening, differently from the case of the evolution by mean curvature flow.

We now introduce the mathematical setting in which we work. Given an initial set $E_0 \subset \mathbb{R}^2$, we define its evolution E_t for $t > 0$ according to a nonlocal curvature flow as follows: the velocity at a point $x \in \partial E_t$ is given by

$$(1.1) \quad \partial_t x \cdot \nu = -H_{E_t}^K(x)$$

where ν is the outer normal at ∂E_t in x . The quantity $H_E^K(x)$ is the K -curvature of E at x , which is defined in the forthcoming formula (1.4). More precisely, we take a function $K : \mathbb{R}^2 \setminus \{0\} \rightarrow [0, +\infty)$ which is a rotationally invariant kernel, namely

$$(1.2) \quad K(x) = K_0(|x|),$$

for some $K_0 : (0, +\infty) \rightarrow [0, +\infty)$. We assume that

$$(1.3) \quad \min\{1, |x|\} K(x) \in L^1(\mathbb{R}^2), \quad \text{i.e.} \quad \int_0^1 \rho^2 K_0(\rho) d\rho + \int_1^{+\infty} \rho K_0(\rho) d\rho < +\infty.$$

Given $E \subset \mathbb{R}^2$ and $x \in \partial E$ we define the K -curvature of E at x , defined by

$$(1.4) \quad H_E^K(x) := \lim_{\varepsilon \searrow 0} \int_{\mathbb{R}^2 \setminus B_\varepsilon(x)} \left(\chi_{\mathbb{R}^2 \setminus E}(y) - \chi_E(y) \right) K(x-y) dy,$$

where, as usual,

$$\chi_E(y) := \begin{cases} 1 & \text{if } y \in E, \\ 0 & \text{if } y \notin E. \end{cases}$$

We point out that (1.2) is a very mild integrability assumption, compatible with the structure of nonlocal minimal surfaces (see e.g. condition (1.5) in [11]) and which fits the requirements in [8, 16] in order to have existence and uniqueness for the level set flow associated to (1.1) (see Appendix A for the details about this matter).

Furthermore, when $K(x) = \frac{1}{|x|^{2+s}}$ for some $s \in (0, 1)$, we will denote the K -curvature of a set E at a point x as $H_E^s(x)$, and we indicate it as the fractional mean curvature of the set E at x .

While the setting in (1.4) makes clear sense for sets with $C^{1,1}$ -boundaries, as customary we also use the notion of K -curvatures for sets which are locally the graphs of continuous functions: in this case, the K -curvature may be also infinite and the definition is in the sense of viscosity (see [8, 16] and Section 5 in [6]).

We observe that the curvature defined in (1.4) is the the first variation of the following nonlocal perimeter functional, see [7, 17],

$$(1.5) \quad \text{Per}_K(E) := \int_E \int_{\mathbb{R}^2 \setminus E} K(x-y) dx dy,$$

and so the geometric evolution law in (1.1) can be interpreted as the L^2 gradient flow of this perimeter functional, as proved in [8].

The existence and uniqueness of solutions for the K -curvature flow in (1.1) in the viscosity sense have been investigated in [16] by introducing the level set formulation of the geometric evolution problem (1.1) and a proper notion of viscosity solution. We refer to [8] for a general framework for the analysis via the level set formulation of a wide class of local and nonlocal translation-invariant geometric flows.

The level set flow associated to (1.1) can be defined as follows. Given an initial set $E \subset \mathbb{R}^2$ and $C := \partial E$, we choose a bounded Lipschitz continuous function $u_E : \mathbb{R}^2 \rightarrow \mathbb{R}$ such that

$$\begin{aligned} C &= \{x \in \mathbb{R}^2 \text{ s.t. } u_E(x) = 0\} = \partial\{x \in \mathbb{R}^2 \text{ s.t. } u_E(x) \geq 0\} \\ \text{and } E &= \{x \in \mathbb{R}^2 \text{ s.t. } u_E(x) \geq 0\}. \end{aligned}$$

Let also $u_E(x, t)$ be the viscosity solution of the following nonlocal parabolic problem

$$(1.6) \quad \begin{cases} \partial_t u(x, t) + |Du(x, t)| H_{\{y|u(y,t) \geq u(x,t)\}}^K(x) = 0, \\ u(x, 0) = u_E(x). \end{cases}$$

Then the level set flow of C is given by

$$(1.7) \quad \Sigma_E(t) := \{x \in \mathbb{R}^2 \text{ s.t. } u_E(x, t) = 0\}.$$

We associate to this level set the outer and inner flows defined as follows:

$$(1.8) \quad E^+(t) := \{x \in \mathbb{R}^2 \text{ s.t. } u_E(x, t) \geq 0\} \quad \text{and} \quad E^-(t) := \{x \in \mathbb{R}^2 \text{ s.t. } u_E(x, t) > 0\}.$$

We observe that the equation in (1.6) is geometric, so if we replace the initial condition with any function u_0 with the same level sets $\{u_0 \geq 0\}$ and $\{u_0 > 0\}$, the evolutions $E^+(t)$ and $E^-(t)$ remain the same. For more details, we refer to Appendix A.

The K -curvature flow has been recently studied from different perspectives, in particular the case fractional mean curvature flow, taking into account geometric features such as conservation of the positivity of the fractional mean curvature, conservation of convexity and formation of neckpinch singularities, see [9, 12, 18].

In this paper, we analyze the possible lack of uniqueness for the geometric evolution, i.e. the situation in which $\partial E^+(t) \neq \partial E^-(t)$, in terms of the fattening properties of the zero level set of the viscosity solutions. To this end, we give the following definition:

Definition 1.1. We say that fattening occurs at time $t > 0$ if the set $\Sigma_E(t)$, defined in (1.7), has nonempty interior, i.e.

$$\text{int}(E^+(t) \setminus E^-(t)) \neq \emptyset.$$

We point out that in [9, Section 6], in the case of fractional (anisotropic) mean curvature flow in any dimension, it has been proved that if the initial set $E \subseteq \mathbb{R}^n$ is convex, then the evolution remains convex for all $t > 0$ and $E^+(t) = \overline{E^-(t)}$, so fattening never occurs.

We start with a result about nonfattening of bounded regular sets with positive K -curvature (for the classical case of the mean curvature flow, see [1, 2, 4]).

Theorem 1.2. *Let (1.2) and (1.3) hold. Let $E \subset \mathbb{R}^2$ be a compact set of class $C^{1,1}$ and we assume that there exists $\delta > 0$ such that*

$$(1.9) \quad H_E^K(x) \geq \delta \text{ for every } x \in \partial E.$$

Then $\Sigma_E(t)$ has empty interior for every t .

We point out that, to get the result in Theorem 1.2, the assumption on the regularity of the sets cannot be completely dropped: indeed in the forthcoming Theorem 1.11 we will provide an example of bounded set, with a ‘‘Lipschitz-type’’ singularity and with positive K -curvature, which develops fattening.

1.1. **Evolution of the cross.** We consider now the cross in \mathbb{R}^2 , i.e.

$$(1.10) \quad \mathcal{C} := \{x = (x_1, x_2) \in \mathbb{R}^2 \text{ s.t. } |x_1| \geq |x_2|\}.$$

It is well known, see [13], that the evolution of the cross according to the curvature flow immediately develops fattening for $t > 0$. So, an interesting question is if the same phenomenon appears also for general nonlocal curvature flows as (1.1), for kernels which satisfy (1.2) and (1.3). We show that actually the fattening feature in nonlocal curvature flows is very sensible to the specific properties of the kernel since it depends on the strength of the interactions: we identify in particular two classes of kernels, giving fattening of the cross in the first class, i.e. for kernels which satisfy (1.16), (1.17) below, and nonfattening of the cross in the second class, i.e. for kernels which satisfy (1.22) below.

Remark 1.3. Recalling the notation in (1.7), we observe that

$$(1.11) \quad \{x = (x_1, x_2) \in \mathbb{R}^2 \text{ s.t. } |x_1| = |x_2|\} \subseteq \Sigma_{\mathcal{C}}(t) \quad \text{for all } t > 0.$$

Indeed, up to a rotation of coordinate system, we write $\mathcal{C} = \{(y_1, y_2) \in \mathbb{R}^2 \text{ s.t. } y_1 y_2 \geq 0\}$. Define a bounded Lipschitz function u_0 such that $u_0(y_1, y_2) = u_0(-y_1, -y_2) = -u_0(-y_1, y_2) = -u_0(y_1, -y_2)$, and such that $\mathcal{C} = \{(y_1, y_2) \in \mathbb{R}^2 \text{ s.t. } u_0(y_1, y_2) \geq 0\}$. Then the solution to (1.6) with initial condition u_0 satisfies

$$u(y_1, y_2, t) = u(-y_1, -y_2, t) = -u(-y_1, y_2, t) = -u(y_1, -y_2, t),$$

see Appendix A. In particular this implies that $\{(y_1, y_2) \in \mathbb{R}^2 \text{ s.t. } y_1 y_2 = 0\} \subseteq \{(y_1, y_2) \in \mathbb{R}^2 \text{ s.t. } u(y_1, y_2, t) = 0\} = \Sigma_{\mathcal{C}}(t)$, that is (1.11) once we rotate back.

We introduce the function

$$(1.12) \quad \Psi(r) := \int_{B_{r/4}(7r/4, 0)} K(x) dx.$$

In our framework, the function $\Psi(r)$ plays a crucial role in quantitative K -curvature estimates. Notice that when $K(x) = \frac{1}{|x|^{2+s}}$ with $s \in (0, 1)$, the function $\Psi(r)$ reduces, up to multiplicative constants, to $\frac{1}{r^s}$.

We define, for any $r > 0$, the ‘‘perturbed cross’’

$$(1.13) \quad \mathcal{C}_r := [-r, r]^2 \cup \mathcal{C}.$$

Then, we have:

Proposition 1.4. *Assume that (1.2) and (1.3) hold true. Then, we have that*

$$(1.14) \quad H_{\mathcal{C}_r}^K(p) \leq 0$$

for any $p \in \partial \mathcal{C}_r$. Also, for any $t \in [-r, r]$,

$$(1.15) \quad H_{\mathcal{C}_r}^K(t, r) \leq -2\Psi(r).$$

Proposition 1.4 provides the cornerstone to detect the fattening phenomenon of the K -curvature flow emanating from the cross, when the kernel K satisfies

$$(1.16) \quad \int_0^1 \frac{d\rho}{\Psi(\rho)} < +\infty.$$

We will need also the following technical assumption: there exists $r_0 > 0$ such that for all $r \in (0, r_0)$,

$$(1.17) \quad \inf_{p \in B_{3\sqrt{2}r}} \int_{B_{r/4}(3r/4, 0) - p} K(x) dx > 0.$$

This assumption is trivially satisfied if $K > 0$ in $B_{(3\sqrt{2}+1)r_0}$.

Indeed in this case, we have that, for short times, the set $\Sigma_{\mathcal{C}}(t)$ contains a ball centered at the origin (see¹ Figure 1), according to the following result:

¹The pictures of this paper have just a qualitative and exemplifying purpose, to favor the intuition and make the reading simpler. They are sketchy, not quantitatively accurate and they are not the outcome of any rigorous simulation.




FIGURE 1. The fattening phenomenon described in Theorem 1.5.

Theorem 1.5. Assume that (1.2), (1.3), (1.16) and (1.17) hold true. For $r \in (0, 1)$, we define

$$(1.18) \quad \Lambda(r) := \int_0^r \frac{d\rho}{\Psi(\rho)}.$$

Then, there exists $T > 0$ such that

$$(1.19) \quad B_{r(t)} \subset \Sigma_{\mathfrak{c}}(t)$$

for any $t \in (0, T)$, where $r(t)$ is defined implicitly by

$$(1.20) \quad \Lambda(r(t)) = t.$$

We notice that the setting in (1.18) is well defined in view of the structural assumption in (1.16) and $\Lambda(r)$, as defined in (1.18), is strictly increasing, which makes the implicit definition in (1.20) well posed.

Remark 1.6. We point out that the structural assumptions in (1.3) and (1.16) are satisfied by kernels of the form $K(x) = \frac{1}{|x|^{2+s}}$ for some $s \in (0, 1)$, or more generally by kernels such that

$$(1.21) \quad K \in L^1(\mathbb{R}^2 \setminus B_1) \quad \text{and} \quad \frac{1}{C|x|^\alpha} \leq K(x) \leq \frac{C}{|x|^\beta}, \quad \text{with } \alpha > 1, \beta < 3, C \geq 1, \text{ for any } x \in B_1.$$

Indeed, the upper bound for K in (1.21) plainly implies (1.3). Moreover, the lower bound for K in (1.21) implies that

$$\Psi(r) = \int_{B_{r/4}(7r/4, 0)} K(x) dx \geq \int_{B_{r/4}(7r/4, 0)} \frac{1}{|x|^\alpha} dx \geq \frac{1}{(2r)^\alpha} |B_{r/4}| = C_0 r^{2-\alpha}$$

where $C_0 > 0$ is independent of r , and this yields (1.16). Finally as for (1.17), we observe that it is trivially satisfied.

Note that $r(t)$ defined in (1.20) satisfies $r(t) \geq C_0 t^{\frac{1}{\alpha-1}}$, in particular, in the case $K(x) = \frac{1}{|x|^{2+s}}$, $r(t)$ is proportional to $t^{\frac{1}{1+s}}$.

As a counterpart of Theorem 1.5, we show that the fattening phenomenon does not occur in straight crosses when the interaction kernel has sufficiently strong integrability properties. Namely, we have that:

Theorem 1.7. Assume (1.2) and (1.3). Suppose also that

$$(1.22) \quad \begin{aligned} &K_0 \leq K_1, \text{ with } K_1 \text{ nonincreasing and} \\ &\Phi(r) := \int_{[-r, r] \times \mathbb{R}} K_1(|x|) dx < +\infty, \end{aligned}$$

for any $r > 0$, and that

$$(1.23) \quad \lim_{\delta \searrow 0} \int_{\delta}^1 \frac{d\tau}{\Phi(\tau)} = +\infty.$$

Then

$$(1.24) \quad \text{the evolution of } \mathcal{C} \text{ under the } K\text{-curvature flow coincides with } \mathcal{C} \text{ itself.}$$

Remark 1.8. We notice that conditions (1.3), (1.22) and (1.23) are satisfied by kernels K such that K_0 is nonincreasing, and which satisfy

$$(1.25) \quad K \in L^1(\mathbb{R}^2 \setminus B_1) \quad \text{and} \quad K(x) \leq \frac{C}{|x|^\alpha}, \quad \text{with } \alpha \in (0, 1], C > 0, \text{ for any } x \in B_1.$$

Indeed, we observe first that in this case (1.3) is automatically satisfied. Moreover, from (1.25), we can take $K_1 := K_0$ in (1.22) and have that

$$\begin{aligned} \Phi(r) &= \int_{[-r,r] \times \mathbb{R}} K_0(|x|) dx \\ &\leq \int_{B_r} \frac{C}{|x|^\alpha} dx + \int_{[-r,r]^2 \setminus B_r} K_0(|x|) dx + \int_{[-r,r] \times ((-\infty, -r] \cup [r, +\infty))} K_0(|x|) dx \\ &\leq Cr^{2-\alpha} + 4r \int_r^{+\infty} K_0(x_2) dx_2 \\ &\leq Cr^{2-\alpha} + Cr \left(\int_r^1 \frac{dx_2}{x_2^\alpha} + 1 \right) \\ &\leq Cr |\log r|, \end{aligned}$$

up to renaming $C > 0$, and so (1.23) is satisfied.

We also observe that condition (1.25) is somewhat complementary to (1.21).

1.2. A remark on K -minimal cones. As a byproduct of the results that we discussed in Subsection 1.1, we observe that actually the cross is not a K -minimal set for the K -perimeter in \mathbb{R}^2 , obtaining an alternative (and more general) proof of a result discussed in Proposition 5.2.3 of [5] for the fractional perimeter (see [19] for a full regularity theory of fractional minimal cones in the plane).

For this, we define

$$(1.26) \quad \text{Per}_K(E, B_R) := \int_{E \cap B_R} \int_{\mathbb{R}^2 \setminus E} K(x-y) dx dy + \int_{E \setminus B_R} \int_{B_R \setminus E} K(x-y) dx dy.$$

Then, we say that E is a minimizer for Per_K in the ball B_R if

$$\text{Per}_K(E, B_R) \leq \text{Per}_K(F, B_R)$$

for every measurable set F such that $E \setminus B_R = F \setminus B_R$.


Also, a measurable set $E \subset \mathbb{R}^2$ is said to be K -minimal for the K -perimeter if it is a minimizer for Per_K in every ball B_R . Then, we have:

Proposition 1.9. *Let (1.2) and (1.3) hold, and assume that K is not identically zero. Then \mathcal{C} , as defined in (1.10), is not K -minimal for the K -perimeter.*

1.3. Fractional curvature evolution of starshaped sets. Now we restrict ourselves to the case of homogeneous kernels K , i.e. we consider the case (up to multiplicative constants) in which

$$(1.27) \quad K_0(r) = \frac{1}{r^{2+s}}, \quad \text{with } s \in (0, 1).$$

We start by observing that strictly starshaped sets never fattens, similarly as for the (local) curvature flow (see [20]). A similar result has also been observed in [9, Remark 6.4].

FIGURE 2. The double droplet \mathcal{G} .

Proposition 1.10. *Assume (1.27). Let $\mathbb{S}_1 := \mathbb{R}/(2\pi\mathbb{Z})$ and $f : \mathbb{S}_1 \rightarrow (0, +\infty)$ be a continuous positive function and let $E \subset \mathbb{R}^2$ such that*

$$(1.28) \quad E = \{0\} \cup \left\{ x \in \mathbb{R}^2 \text{ s.t. } x \neq 0, |x| \leq f\left(\frac{x}{|x|}\right) \right\}.$$

Then, the set $\Sigma_E(t)$ has empty interior for all $t > 0$.

Then we show that in general, for starshaped sets E which do not satisfy (1.28), we can expect either fattening or nonfattening. We provide two different examples of such sets, which are particularly interesting in our opinion, since they model two different type of singularities that can arise in the geometric evolution of closed curves in \mathbb{R}^2 , that is the ‘‘Lipschitz-type’’ singularity, and the ‘‘cusp singularity’’. The first example is the ‘‘double droplet’’ in Figure 2, namely

$$(1.29) \quad \mathcal{G} := \mathcal{G}_+ \cup \mathcal{G}_-,$$

where \mathcal{G}_+ is the convex hull of $B_1(-1, 1)$ with the origin, and \mathcal{G}_- the convex hull of $B_1(1, -1)$ with the origin. The second example is given by two tangent balls

$$(1.30) \quad \mathcal{O} := B_1(-1, 0) \cup B_1(1, 0).$$

We prove that fattening phenomenon occurs in the first case, whereas it does not occur in the second. It is also interesting to observe that the evolution of \mathcal{O} by curvature flow immediately develops fattening, see [3].

We start by considering the evolution of the set \mathcal{G} defined in (1.29). Note that this provides an example of bounded set with positive K -curvature (being contained in a cross with zero K -curvature), whose evolution develops fattening near the origin, as sketched in Figure 3 and detailed in the following statement.

Theorem 1.11. *Assume (1.27). Then there exist $\hat{c}, T > 0$ such that*

$$(1.31) \quad B_{r(t)} \subset \Sigma_{\mathcal{G}}(t)$$

for any $t \in (0, T)$, where

$$(1.32) \quad r(t) := \hat{c}t^{1/(1+s)}.$$

Remark 1.12. The same result as in Theorem 1.11 holds more generally for kernels K_0 which satisfy (1.2), (1.3), (1.16) and

$$(1.33) \quad \frac{a}{r^{2+s}} \leq K_0(r) \leq \frac{\bar{a}}{r^{2+s}} \quad \text{for all } r > 0$$

for some suitable $\bar{a} \geq a > 0$.

We now consider the case of two tangent balls as in (1.30), and we show that $\mathcal{O}(t)$ presents no fattening phenomenon, according to the statement below.

Theorem 1.13. *Assume (1.27). Then the set $\Sigma_{\mathcal{O}}(t)$ has empty interior for all $t > 0$.*




FIGURE 3. *The fattening phenomenon described in Theorem 1.11.*




FIGURE 4. *The evolution of two tangent balls described in Theorem 1.13.*

The evolution of the double ball is sketched in Figure 4: roughly speaking, the set shrinks at its surroundings, emanating some mass from the origin, but it does not possess “gray regions” at its boundary.

The rest of the paper is organized as follows. Section 2 deals with the fact that the evolution starting from regular sets with positive K -curvature does not fatten and it contains the proof of Theorem 1.2. In Section 3 we prove Proposition 1.4 and the fattening of the evolution starting from the cross, under assumption (1.16), as stated in Theorem 1.5.

In Section 4, we show that under assumption (1.22) the evolution starting from the cross does not fatten, but coincides with the cross itself, that is we prove Theorem 1.7.

Section 5 contains the proof of the fact that the cross is never a K -minimal set for Per_K , thus establishing Proposition 1.9.

The last three sections present the evolution under the fractional curvature flow, i.e., we assume that $K(x) = \frac{1}{|x|^{2+s}}$. In particular, Section 6 is devoted to the proof of the fact that the fractional curvature evolution of strictly starshaped sets does not present fattening, which gives Proposition 1.10.

In Section 7 we show an example of a compact set with positive K -curvature, that is the double droplet, whose fractional curvature evolution presents fattening, thus proving Theorem 1.11.

Then, in Section 8 we show that the fractional curvature evolution starting from two tangent balls does not fatten, which establishes Theorem 1.13.

In Appendix A we review some basic facts about level set flow, moreover we provide some auxiliary results about comparison with geometric barriers and other basic properties of the evolution which are exploited in the proofs of the main results.

Notation. We denote by $B_r \subset \mathbb{R}^2$ the ball centered at $(0,0)$ of radius r , and by $B_r(x_1, x_2)$ the ball of radius r and center $x = (x_1, x_2) \in \mathbb{R}^2$.

Moreover $e_1 = (1, 0)$ and $e_2 = (0, 1)$ and $\mathbb{S}_1 = \mathbb{R}/(2\pi\mathbb{Z}) = \{e \in \mathbb{R}^2 \text{ s.t. } |e| = 1\}$.

For a given closed set E , and for any $x \in \mathbb{R}^2 \setminus E$ we denote by $\text{dist}(x, E)$ the distance from x to E , that is

$$\text{dist}(x, E) := \inf_{y \in E} |x - y|.$$

Moreover, we will denote with $d_E(x)$ the signed distance function to $C = \partial E$, with the sign convention of being positive inside E and negative outside, that is

$$(1.34) \quad d_E(x) = \begin{cases} \text{dist}(x, \mathbb{R}^2 \setminus E) & \text{if } x \in \overline{E}, \\ -\text{dist}(x, E) & \text{if } x \in \mathbb{R}^2 \setminus E. \end{cases}$$

Finally, given two sets $E, F \subset \mathbb{R}^2$, we denote by $d(E, F)$ the distance between the boundary of E and the boundary of F , that is

$$(1.35) \quad d(E, F) := \min_{\substack{x \in \partial E \\ y \in \partial F}} |x - y|.$$

2. REGULAR SETS OF POSITIVE K -CURVATURE AND PROOF OF THEOREM 1.2

Proof of Theorem 1.2. We recall the continuity in $C^{1,1}$ of the K -curvature proved in [8]. Namely, if E^ε is a family of compact sets with boundaries in $C^{1,1}$ such that $E^\varepsilon \rightarrow E$ in $C^{1,1}$ (in the sense that the boundaries converges in $C^{1,1}$) and $x^\varepsilon \in \partial E^\varepsilon \rightarrow x \in \partial E$, then $H_{E^\varepsilon}^K(x^\varepsilon) \rightarrow H_E^K(x)$, as $\varepsilon \searrow 0$.

Now, let E be as in the statement of Theorem 1.2, and define, for $r > 0$,

$$E^r := \{x \in \mathbb{R}^2 \text{ s.t. } d_E(x) \geq -r\}.$$

Then, using also (1.9), we find that there exists $\varepsilon_0 > 0$ such that, for all $\varepsilon \in (0, \varepsilon_0)$, there exists $0 < \delta(\varepsilon) \leq \delta$ such that

$$\min_{x \in \partial E^\varepsilon} H_{E^\varepsilon}^K(x) \geq \delta(\varepsilon) > 0.$$

Fix $\varepsilon < \varepsilon_0$ and let $\bar{\delta} := \inf_{h \in [\varepsilon/2, \varepsilon]} \delta(h) > 0$. For all $t \in [0, \frac{\varepsilon}{\bar{\delta}}]$ we define

$$C(t) := E^{\varepsilon - \frac{\bar{\delta}}{2}t}.$$

We observe that $C(t)$ is a supersolution to (1.1), in the sense that it satisfies (A.6). Indeed,

$$\partial_t x \cdot \nu = -\frac{\bar{\delta}}{2} \geq -H_{C(t)}^K(x) + \delta,$$

with any $\delta < \bar{\delta}/2$.

Since $E \subseteq E^\varepsilon = C(0)$, by Proposition A.10, we get that

$$E^+(s) \subseteq C(s) = E^{\varepsilon - \frac{\bar{\delta}}{2}s}$$

and $d\left(E^+(s), E^{\varepsilon - \frac{\bar{\delta}}{2}s}\right) \geq \varepsilon,$

for all $s \in [0, \frac{\varepsilon}{\bar{\delta}}]$. This gives that, if

$$u_E(x) := \max\{-M, \min\{M, d_E(x)\}\},$$


and $M > 0$ is sufficiently large, then

$$E^+(s) \subseteq \left\{x \in \mathbb{R}^2 \text{ s.t. } u_E(x) \geq \frac{\bar{\delta}}{4}s\right\} \quad \text{for all } s \in \left[0, \frac{\varepsilon}{\bar{\delta}}\right],$$

with $d\left(E^+(s), \left\{x \in \mathbb{R}^2 \text{ s.t. } u_E(x) \geq \frac{\bar{\delta}}{4}s\right\}\right) \geq \frac{\bar{\delta}}{4}s.$

Then, by the Comparison Principle in Corollary A.8, we get that

$$E^+(t+s) \subseteq \left\{x \in \mathbb{R}^2 \text{ s.t. } u_E(x, t) > \frac{\bar{\delta}}{4}s\right\},$$

FIGURE 5. *The set \mathcal{D}_r .*

for all $t > 0$. This implies immediately that

$$E^+(t+s) \subseteq E^-(t), \quad \text{for all } t > 0, s \in \left(0, \frac{\varepsilon}{\delta}\right].$$

Therefore, recalling Proposition A.12, we get

$$|\text{int}(E^+(t)) \setminus \overline{E^-(t)}| \leq \limsup_{s \searrow 0} |\text{int}(E^+(t))| - |E^+(t+s)| = |\text{int}(E^+(t))| - \liminf_{s \searrow 0} |E^+(t+s)| \leq 0.$$

This gives the desired statement in Theorem 1.2. \square

3. K -CURVATURE OF THE PERTURBED CROSS AND PROOFS OF PROPOSITION 1.4 AND OF THEOREM 1.5

Here, we consider the cross \mathcal{C} introduced in (1.10) and the perturbed cross defined in (1.13). We also make use of the notation in (1.12). Then, we have:

Lemma 3.1. *Assume that (1.2) and (1.3) hold true. Then, for any $t \in [-r, r]$,*

$$H_{\mathcal{C}_r}^K(t, r) \leq -2\Psi(r).$$

Proof. Let

$$\mathcal{T}_r := \left((-r, r)^2 \setminus \mathcal{C} \right) \cap \{x_2 < 0\}$$

and

$$\mathcal{D}_r := \mathcal{C}_r \setminus \mathcal{T}_r,$$

see Figure 5. Notice that \mathcal{C}_r is the disjoint union of \mathcal{D}_r and \mathcal{T}_r , hence

$$\chi_{\mathcal{C}_r} = \chi_{\mathcal{D}_r} + \chi_{\mathcal{T}_r},$$

while $\mathbb{R}^2 \setminus \mathcal{D}_r$ is the disjoint union of $\mathbb{R}^2 \setminus \mathcal{C}_r$ and \mathcal{T}_r , which gives that

$$\chi_{\mathbb{R}^2 \setminus \mathcal{D}_r} = \chi_{\mathbb{R}^2 \setminus \mathcal{C}_r} + \chi_{\mathcal{T}_r}.$$

Hence, we find that

$$(3.1) \quad \chi_{\mathbb{R}^2 \setminus \mathcal{C}_r} - \chi_{\mathcal{C}_r} = \chi_{\mathbb{R}^2 \setminus \mathcal{D}_r} - \chi_{\mathcal{D}_r} - 2\chi_{\mathcal{T}_r}.$$

Now, we claim that, for any $t \in [-r, r]$,

$$(3.2) \quad H_{\mathcal{D}_r}^K(t, r) \leq 0.$$




FIGURE 6. *Splitting the set \mathcal{D}_r and its complement into isometric regions.*

To this end, we partition \mathbb{R}^2 into different regions, as depicted in Figure 6, and we use the notation, for each set $Y \subseteq \mathbb{R}^2$,

$$(3.3) \quad \mathcal{H}(Y) := \lim_{\varepsilon \searrow 0} \int_{Y \setminus B_\varepsilon(t,r)} K(x - (t,r)) dx.$$

In this way, we can write (1.4) as

$$(3.4) \quad H_{\mathcal{D}_r}^K(t,r) = \mathcal{H}(C) + \mathcal{H}(D) + \mathcal{H}(U') + \mathcal{H}(V') + \mathcal{H}(W') - \mathcal{H}(A) - \mathcal{H}(B) - \mathcal{H}(U) - \mathcal{H}(V) - \mathcal{H}(W).$$

On the other hand, we can use symmetric reflections across the horizontal straight line passing through the pole (t,r) to conclude that $\mathcal{H}(U) = \mathcal{H}(U')$. Similarly, we see that $\mathcal{H}(V) = \mathcal{H}(V')$ and $\mathcal{H}(W) = \mathcal{H}(W')$. As a consequence, the identity in (3.4) becomes

$$(3.5) \quad H_{\mathcal{D}_r}^K(t,r) = \mathcal{H}(C) + \mathcal{H}(D) - \mathcal{H}(A) - \mathcal{H}(B).$$

Now we consider the straight line $\ell := \{x_2 = x_1 - t + r\}$. Notice that ℓ passes through the point (t,r) and it is parallel to two edges of the cross \mathcal{C}_r . Considering the framework in Figure 6, reflecting the set D across ℓ we obtain a set $D' \subseteq B$, and we write $B = D' \cup E$, for a suitable slab E . Similarly, we reflect the set A across ℓ to obtain a set A' which is contained in C , and we write $C = A' \cup F$, for a suitable slab F , see Figure 7.

In further details, if $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is the reflection across ℓ , we have that $T(t,r) = (t,r)$ and $|T(x - (t,r))| = |T(x) - (t,r)| = |x - (t,r)|$ for every $x \in \mathbb{R}^2$, and thus, by (1.2),

$$K(x - (t,r)) = K_0(|x - (t,r)|) = K_0(|T(x - (t,r))|) = K(T(x - (t,r))).$$

Accordingly, since $D = T(D')$,

$$(3.6) \quad \begin{aligned} \mathcal{H}(B) - \mathcal{H}(D) &= \int_B K(x - (t,r)) dx - \int_{T(D')} K(x - (t,r)) dx \\ &= \int_B K(x - (t,r)) dx - \int_{D'} K(x - (t,r)) dx = \int_E K(x - (t,r)) dx, \end{aligned}$$

and similarly

$$(3.7) \quad \mathcal{H}(C) - \mathcal{H}(A) = \int_F K(x - (t,r)) dx.$$

Now we consider the straight line $\ell' := \{x_2 = -x_1 + t + r\}$. Notice that ℓ' passes through the point (t,r) and it is perpendicular to ℓ . We let E' be the reflection across ℓ' of the set E and we notice that $E' \supseteq F$. Therefore

$$\int_E K(x - (t,r)) dx = \int_{E'} K(x - (t,r)) dx \geq \int_F K(x - (t,r)) dx.$$




FIGURE 7. Reflecting D and A across ℓ , being $E := B \setminus D'$ and $F := C \setminus A'$.

From this, (3.6) and (3.7), we obtain

$$\mathcal{H}(C) + \mathcal{H}(D) - \mathcal{H}(A) - \mathcal{H}(B) = \int_F K(x - (t, r)) dx - \int_E K(x - (t, r)) dx \leq 0.$$

This and (3.5) imply the desired result in (3.2).

Then, by (3.1) and (3.2),

$$H_{\mathcal{C}_r}^K(t, r) = H_{\mathcal{D}_r}^K(t, r) - 2 \int_{\mathcal{J}_r} K(y - (t, r)) dy \leq 0 - 2\Psi(r),$$

and this gives the desired result. \square

With this, we are now in the position of completing the proof of Proposition 1.4 via the following argument:

Proof of Proposition 1.4. The claim in (1.15) follows from Lemma 3.1. In addition, we have that $\mathcal{C} \subset \mathcal{C}_r$, due to (1.13). We also observe that if $p \in (\partial\mathcal{C}_r) \setminus [-r, r]^2$, then $p \in \partial\mathcal{C}$. Consequently, by (1.4), for any $p \in (\partial\mathcal{C}_r) \setminus [-r, r]^2$, we have that

$$(3.8) \quad H_{\mathcal{C}}^K(p) \geq H_{\mathcal{C}_r}^K(p).$$

Also, by symmetry, we see that $H_{\mathcal{C}}^K(p) = 0$ at any point $p \in \partial\mathcal{C}$, hence (3.8) gives that $H_{\mathcal{C}_r}^K(p) \leq 0$ for any $p \in (\partial\mathcal{C}_r) \setminus [-r, r]^2$. Since this inequality is also valid when $p \in (\partial\mathcal{C}_r) \cap [-r, r]^2$, due to (1.15), the proof of (1.14) is complete. \square

With Proposition 1.4, we can now construct inner and outer barriers as in Corollary A.11 to complete the proof of Theorem 1.5. This auxiliary construction goes as follows.

Lemma 3.2. *Let \mathcal{C}_r be as in (1.13). Let $R := 3\sqrt{2}r$ and define, for $\lambda \in [0, \frac{r}{2})$,*

$$(3.9) \quad \mathcal{C}_r^\lambda := \{x \in \mathbb{R}^2 \text{ s.t. } d_{\mathcal{C}_r}(x) \leq -\lambda\}.$$

Then, for any $p \in (\partial\mathcal{C}_r^\lambda) \setminus B_R$, we have that $H_{\mathcal{C}_r^\lambda}^K(p) \leq 0$.

Proof. We observe that if $p \in (\partial\mathcal{C}_r^\lambda) \setminus B_R$, then $\partial\mathcal{C}_r^\lambda$ in the vicinity of p is a segment, and there exists a vertical translation of \mathcal{C} by a vector $v_0 := \pm\sqrt{\lambda}e_2$ such that $p \in \mathcal{C} + v_0$ and $\mathcal{C} + v_0 \subset \mathcal{C}_r^\lambda$, see Figure 8. From this, we find that

$$H_{\mathcal{C}_r^\lambda}^K(p) \leq H_{\mathcal{C}+v_0}^K(p) = H_{\mathcal{C}}^K(p - v_0) = 0,$$

as desired. \square




FIGURE 8. The set \mathcal{C}_r^λ , touched from inside at a boundary point by a translation of \mathcal{C} .

With this, we are ready to complete the proof of Theorem 1.5, by arguing as follows.

Proof of Theorem 1.5. The proof is based on the construction of suitable families of geometric sub and super-solutions starting from the perturbed cross \mathcal{C}_r , as defined in (1.13), to which apply Corollary A.11.

We observe that

$$\mathcal{C} = \bigcap_{r>0} \mathcal{C}_r.$$

Moreover, we see that

$$d_{\mathcal{C}}(x) \leq d_{\mathcal{C}_r}(x) \leq d_{\mathcal{C}}(x) + r.$$

These observations, together with the Comparison Principle in Theorem A.5 and Remark A.6, imply that

$$(3.10) \quad \mathcal{C}^+(t) = \bigcap_{r>0} \mathcal{C}_r^+(t), \quad \text{for all } t > 0.$$

Analogously, one can define

$$(3.11) \quad \mathcal{C}^r := (\mathbb{R}^2 \setminus \mathcal{C}) \cup [-r, r]^2.$$

Let Ψ as defined in (1.12). Fixed $r \in (0, r_0)$, where r_0 is as in (1.17), we define $r_*(t)$ to be the solution to the ODE

$$(3.12) \quad \dot{r}_*(t) = \Psi(r_*(t))$$

with initial datum $r_*(0) = r$. We fix $T > 0$ such that $r_*(t) < r_0$ for all $t \in [0, T]$. Recalling the definition of Λ in (1.18), it is easy to check that

$$(3.13) \quad \Lambda(r_*(t)) = t + \Lambda(r), \quad \text{for all } t \in (0, T].$$

Now, by (1.20) and (3.13), we see that

$$(3.14) \quad \Lambda(r_*(t)) = \Lambda(r(t)) + \Lambda(r) \geq \Lambda(r(t)).$$

Now, recalling the setting in (3.9), we take into account the sets $\mathcal{C}_{r_*(t)}$ and $\mathcal{C}_{r_*(t)}^\lambda$, with $\lambda \in [0, \frac{\pi}{2})$ and $t \in [0, T]$, and we claim that these sets satisfy the assumptions in Corollary A.11, item ii). To this end, we observe that, in the vicinity of the angular points of \mathcal{C}_r , the complement of \mathcal{C}_r is a convex set, and therefore condition (A.9) is satisfied by $\mathcal{C}_{r_*(t)}$. Also, we take

$$\delta_1 := \inf_{t \in [0, T]} \Psi(r_*(t)), \quad \delta_2 := \inf_{t \in [0, T]} \inf_{p \in B_{3\sqrt{2}r_*(t)}} \int_{B_{r_*(t)/4}(3r_*(t)/4, 0) - p} K(y) dy \quad \text{and} \quad \delta := \min\{\delta_1, \delta_2\}.$$

Notice that $\delta > 0$ thanks to (1.16) and (1.17). Then, by Proposition 1.4 and (3.12), we get that at any point $x = (x_1, x_2)$ of $\partial\mathcal{C}_{r_*(t)}$ with $x_2 = \pm r_*(t)$, we have that

$$(3.15) \quad -H_{\mathcal{C}_{r_*(t)}}^K(x) \geq 2\Psi(r_*(t)) = \dot{r}_*(t) + \Psi(r_*(t)) \geq \dot{r}_*(t) + \delta_1 \geq \partial_t x \cdot \nu(x) + \delta.$$

In addition, if $x = (x_1, x_2) \in (\partial\mathcal{C}_{r_*(t)}) \cap B_{4R}$ and $|x_2| > r_*(t)$, we have that

$$-H_{\mathcal{C}_{r_*(t)}}^K(x) \geq -H_{\mathcal{C}}^K(x) + \int_{B_{r_*(t)/4}(3r_*(t)/4, 0)} K(y-x) dy = \int_{B_{r_*(t)/4}(3r_*(t)/4, 0)-x} K(y) dy \geq \delta_2 \geq \delta = \partial_t x \cdot \nu(x) + \delta.$$

This and (3.15) give that condition (A.8) is fulfilled by $\mathcal{C}_{r_*(t)}$.

Furthermore, in light of Lemma 3.2, we know that, for any $x \in (\partial\mathcal{C}_{r_*(t)}^\lambda) \setminus B_R$,

$$H_{\mathcal{C}_{r_*(t)}^\lambda}^K(p) \leq 0 = \partial_t x \cdot \nu(x),$$

which says that condition (A.15) is fulfilled by $\mathcal{C}_{r_*(t)}^\lambda$.

Therefore, we are in the position of using Corollary A.11, item ii). In this way, we find that

$$\mathcal{C}_{r_*(t)} \subseteq \mathcal{C}_r^+(t), \quad \text{for all } t \in [0, T].$$

Hence, recalling (3.14),

$$\mathcal{C}_{r(t)} \subseteq \mathcal{C}_r^+(t), \quad \text{for all } t \in [0, T].$$

Taking intersections, in view of (3.10), we obtain that

$$(3.16) \quad \mathcal{C}_{r(t)} \subseteq \mathcal{C}^+(t), \quad \text{for all } t \in [0, T].$$

Analogously, one can use the setting in (3.11), combined with Corollary A.11, item i), and deduce that

$$(3.17) \quad \mathcal{C}^{r(t)} \subseteq (\mathbb{R}^2 \setminus \mathcal{C})^+(t) \quad \text{for all } t \in [0, T].$$

By (3.16) and (3.17) we get

$$[-r(t), r(t)]^2 = \mathcal{C}_{r(t)} \cap \mathcal{C}^{r(t)} \subseteq \mathcal{C}^+(t) \cap (\mathbb{R}^2 \setminus \mathcal{C})^+(t) = \Sigma_{\mathcal{C}}(t),$$

which implies (1.19), as desired. \square

4. MOVING BOXES, WEAK INTERACTION KERNELS AND PROOF OF THEOREM 1.7

To simplify some computation, in this section we operate a rotation of coordinates so that

$$(4.1) \quad \mathcal{C} = \{x \in \mathbb{R}^2 \text{ s.t. } x_1 x_2 \geq 0\}.$$

To prove Theorem 1.7, it is convenient to consider “expanding boxes” built by the following sets. For any $r \in (0, 1)$, we define

$$(4.2) \quad \mathcal{N}_r := \left([r, +\infty) \times [r, +\infty)\right) \cup \left((-\infty, -r] \times (-\infty, -r]\right),$$

see Figure 9.

Then, recalling the notation in (1.22), we have:

Lemma 4.1. *Assume that K satisfies (1.2), (1.3) and (1.22). Then, for any $p \in \partial\mathcal{N}_r$,*



$$H_{\mathcal{N}_r}^K(p) \leq 2\Phi(2r).$$

Proof. We denote by A and B the two connected components of \mathcal{N}_r and consider the straight line ℓ passing through p and tangent to \mathcal{N}_r at p : see Figure 10. By reflection across ℓ , we can consider the regions A' and B' which are symmetric to A and B , respectively. In particular, if $p = (p_1, p_2)$ and $M(x_1, x_2) := (2p_1 - x_1, x_2)$, we have that $M(A \cup B) = A' \cup B'$ and $M(B_\varepsilon(p)) = B_\varepsilon(p)$, and therefore

$$\begin{aligned} \int_{(A' \cup B') \setminus B_\varepsilon(p)} K(p-y) dy &= \int_{M((A \cup B) \setminus B_\varepsilon(p))} K(p-y) dy = \int_{M((A \cup B) \setminus B_\varepsilon(p))} K(p-Mx) dx \\ &= \int_{M((A \cup B) \setminus B_\varepsilon(p))} K(-p_1 + x_1, p_2 - x_2) dx = \int_{(A' \cup B') \setminus B_\varepsilon(p)} K(p-x) dx, \end{aligned}$$

thanks to (1.2). Then, denoting by

$$T := (\mathbb{R}^2 \setminus \mathcal{N}_r) \setminus (A' \cup B'),$$

FIGURE 9. *The set \mathcal{N}_r .*FIGURE 10. *Simplifications in the computations of Lemma 4.1.*

which is the “white region” in Figure 10, we see that

$$(4.3) \quad \begin{aligned} H_{\mathcal{N}_r}^K(p) &= \lim_{\varepsilon \searrow 0} \int_{(A' \cup B') \setminus B_\varepsilon(p)} K(p-x) dx - \int_{(A \cup B) \setminus B_\varepsilon(p)} K(p-x) dx + \int_T K(p-x) dx \\ &= \int_T K(p-x) dx. \end{aligned}$$

Up to rotations, we may assume that

$$(4.4) \quad T = (\mathbb{R} \times [-r, r]) \cup ([-r, 3r] \times (-\infty, -r]).$$

Recalling (1.22), and that $p_1 = r$, we get

$$(4.5) \quad \int_{[-r, 3r] \times (-\infty, -r]} K(x-p) dx \leq \int_{[-r, 3r] \times (-\infty, -r]} K_1(|x-p|) dx \leq \int_{[-r, 3r] \times \mathbb{R}} K_1(|x-p|) dx = \Phi(2r)$$




FIGURE 11. The set \mathcal{N}_r^λ , touched from inside at a boundary point by a translation of \mathcal{N}_r .

where Φ is defined in (1.22). Moreover, since $p_1 = r$ and $p_2 \geq r$, and K_1 is nonincreasing, we get that $K_1(|x - p|) \leq K_1(|x - (r, r)|)$, for every $x \in \mathbb{R} \times [-r, r]$. As consequence,

$$\begin{aligned} \int_{\mathbb{R} \times [-r, r]} K(x - p) dx &\leq \int_{\mathbb{R} \times [-r, r]} K_1(|x - p|) dx \\ &\leq \int_{\mathbb{R} \times [-r, r]} K_1(|x - (r, r)|) dx \leq \int_{\mathbb{R} \times [-r, 3r]} K_1(|x - (r, r)|) dx = \Phi(2r). \end{aligned}$$

From this and (4.5), and recalling (4.4), we obtain that

$$\int_T K(p - x) dx \leq 2\Phi(2r).$$

This and (4.3) give the desired result. \square

For $\lambda \in (0, r)$ we define the sets

$$(4.6) \quad \mathcal{N}_r^\lambda := \{x \in \mathbb{R}^2 \text{ s.t. } d_{\mathcal{N}_r}(x) \geq -\lambda\}.$$

We observe that for any $x \in \partial\mathcal{N}_r^\lambda$ there exists a unique point $x' \in \partial\mathcal{N}_r$ such that $|x - x'| = d(\mathcal{N}_r^\lambda, \mathcal{N}_r) = \lambda$. Letting $v_x := x - x'$, it follows that $\mathcal{N}_r + v_x \subset \mathcal{N}_r^\lambda$, see Figure 11. This and Lemma 4.1 give that

$$(4.7) \quad H_{\mathcal{N}_r^\lambda}^K(x) \leq H_{\mathcal{N}_r}^K(x + v_x) \leq 2\Phi(2r) \quad \text{for any } x \in \partial\mathcal{N}_r^\lambda.$$

With this preliminary work, we can prove Theorem 1.7.

Proof of Theorem 1.7. We note that $\mathcal{M}_r := \mathcal{N}_r^{r/2} \subseteq \mathcal{C}$, being \mathcal{C} defined in (4.1) and $\mathcal{N}_r^{r/2}$ defined in (4.6), with $\lambda = r/2$. Moreover, we have that $d(\mathcal{C}, \mathcal{M}_r) = r/2 > 0$. Hence, by Corollary A.8 we get that $\mathcal{M}_r^+(t) \subseteq \mathcal{C}^-(t)$ for all $t > 0$. In particular, since

$$\bigcup_{r>0} \mathcal{M}_r = \text{int}(\mathcal{C}),$$

we see that

$$(4.8) \quad \bigcup_{r>0} \mathcal{M}_r^+(t) = \mathcal{C}^-(t).$$

Our aim is to construct starting from \mathcal{M}_r a continuous family of geometric subsolutions and then apply Proposition A.10. Fixed $\varrho \in (0, 1)$, we define

$$F_\varrho(r) := \int_\varrho^r \frac{d\vartheta}{6\Phi(2\vartheta)}.$$

Notice that F_ϱ is strictly increasing, so we can consider its inverse G_ϱ in such a way that $F_\varrho(G_\varrho(t)) = t$. Then, for $t \in [0, T]$, we set $r_\varrho(t) := G_\varrho(t)$ and we consider the evolving sets $\mathcal{M}_{r_\varrho(t)}$. We remark that

$$F_\varrho(\varrho) = 0 = F_\varrho(G_\varrho(0)) = F_\varrho(r_\varrho(0)),$$

and so $r_\varrho(0) = \varrho$. In addition, the outer normal velocity of $\mathcal{M}_{r_\varrho(t)}$ is

$$(4.9) \quad -\dot{r}_\varrho(t) + \frac{1}{2}\dot{r}_\varrho(t) = -\frac{1}{2}G'_\varrho(t) = -\frac{1}{2F'_\varrho(G_\varrho(t))} = -3\Phi(2G_\varrho(t)) = -3\Phi(2r_\varrho(t)).$$

So, if

$$\delta := \Phi(2\rho) = \min_{r \in [\varrho, r_\varrho(T)]} \Phi(2r),$$

we have that

$$(4.10) \quad \partial_t x \cdot \nu(x) = -\frac{1}{2} \dot{r}_\varrho(t) = -2\Phi(2r_\varrho(t)) - \Phi(2r_\varrho(t)) \leq -H_{\mathcal{M}_{r_\varrho(t)}}^K(x) - \delta$$

for all $x \in \partial\mathcal{M}_{r_\varrho(t)}$, thanks to (4.7).

We observe that (4.10) says that (A.8) is satisfied by $\mathcal{M}_{r_\varrho(t)}$. So, to exploit Corollary A.11, we now want to check that condition (A.15) is satisfied by the set

$$\mathcal{M}_{r_\varrho(t)}^\lambda := \{x \in \mathbb{R}^2 \text{ s.t. } d_{\mathcal{M}_{r_\varrho(t)}}(x) \geq -\lambda\} \quad \text{for } \lambda \in (0, \rho).$$

We exploit again the estimate (4.7) which gives that

$$H_{\mathcal{M}_{r_\varrho(t)}^\lambda}^K(x) \leq 2\Phi(2r_\varrho(t)) \quad \text{for any } x \in \partial\mathcal{M}_{r_\varrho(t)}^\lambda.$$

Thus, in view of (4.9),

$$\partial_t x \cdot \nu(x) = -\frac{1}{2} \dot{r}_\varrho(t) = -3\Phi(2r_\varrho(t)) \leq -H_{\mathcal{M}_{r_\varrho(t)}^\lambda}^K(x).$$

This gives that $\mathcal{M}_{r_\varrho(t)}^\lambda$ satisfies condition (A.15) and therefore we can apply Corollary A.11, item ii).

Then, it follows that, for all $\varrho \in (0, 1)$,

$$(4.11) \quad \mathcal{M}_{r_\varrho(t)} \subseteq \mathcal{M}_\varrho^+(t).$$

Also, for any $t > 0$, we claim that

$$(4.12) \quad \lim_{\varrho \searrow 0} r_\varrho(t) = 0.$$

To prove this, we argue by contradiction and suppose that $r_{\varrho_k}(t) \geq a_0$, for some $a_0 > 0$ and some infinitesimal sequence ϱ_k . Then,

$$t = F_{\varrho_k}(G_{\varrho_k}(t)) = F_{\varrho_k}(r_{\varrho_k}(t)) \geq F_{\varrho_k}(a_0) = \int_{\varrho_k}^{a_0} \frac{d\vartheta}{6\Phi(2\vartheta)} = \frac{1}{12} \int_{2\varrho_k}^{2a_0} \frac{d\tau}{\Phi(\tau)}.$$

This is in contradiction with (1.23) and so it proves (4.12).

In view of (4.12), we find that

$$\bigcup_{\varrho > 0} \mathcal{M}_{r_\varrho(t)} = \text{int } \mathcal{C}.$$

So, recalling (4.8) and (4.11), we conclude that

$$(4.13) \quad \text{int } \mathcal{C} = \bigcup_{\varrho > 0} \mathcal{M}_{r_\varrho(t)} \subseteq \bigcup_{\varrho > 0} \mathcal{M}_\varrho^+(t) = \mathcal{C}^-(t) \quad \text{for all } t \in [0, T].$$

Analogously, one can define

$$\mathcal{N}^r := \left((-\infty, -r] \times [r, +\infty) \right) \cup \left([r, +\infty) \times (-\infty, -r] \right), \quad \mathcal{M}^r = (\mathcal{N}^r)^{r/2} := \{x \in \mathbb{R}^2 \text{ s.t. } d_{\mathcal{N}^r}(x) \geq -\lambda\}.$$

and see that

$$(4.14) \quad \text{int } (\mathbb{R}^2 \setminus \mathcal{C}) \subseteq \mathbb{R}^2 \setminus \mathcal{C}^-(t) \quad \text{for all } t \in [0, T].$$

Putting together (4.13) and (4.14), we conclude that

$$\text{int } \mathcal{C} \subseteq \mathcal{C}^-(t) \subseteq \mathcal{C}^+(t) \subseteq \mathcal{C},$$

and so $\Sigma_{\mathcal{C}}(t) = \partial\mathcal{C}$, thus establishing (1.24). \square

5. K -MINIMAL CONES AND PROOF OF PROPOSITION 1.9

In this section we show that \mathcal{C} , as defined in (1.10), is never a K -minimal set, under the assumptions (1.2) and (1.3), namely we prove Proposition 1.9. This will be proved using the family of perturbed crosses \mathcal{C}_r introduced in (1.13) and the fact that H_E^K is the first variation of the nonlocal perimeter Per_K defined in (1.5), as shown in [8].

Proof of Proposition 1.9. With the notation in (1.10) and (1.13), we claim that there exists $r > 0$ such that, for all $R > \sqrt{2}r$,

$$(5.1) \quad \text{Per}_K(\mathcal{C}_r, B_R) < \text{Per}_K(\mathcal{C}, B_R).$$

Let $r > 0$ and $R > \sqrt{2}r$, so that $\mathcal{C}_r \setminus B_R = \mathcal{C} \setminus B_R$. Let

$$W_r := \mathcal{C}_r \setminus \mathcal{C} \subseteq B_R.$$

Let $\delta \in (0, r)$ and $K_\delta(y) := K(y)(1 - \chi_{B_\delta}(y))$. We define $\text{Per}_\delta(E)$ as in (1.5), $\text{Per}_\delta(E, B_R)$ as in (1.26), and H_E^δ as in (1.4), with K_δ in place of K . In this setting, we get that

$$(5.2) \quad \text{Per}_\delta(W_r) = \text{Per}_\delta(W_r, B_R) = \text{Per}_\delta(\mathcal{C}_r, B_R) - \text{Per}_\delta(\mathcal{C}, B_R) + 2 \int_{W_r} \int_{\mathcal{C}} K_\delta(x-y) dx dy.$$

We also observe that

$$\text{Per}_\delta(W_r) = \int_{W_r} \int_{\mathbb{R}^2 \setminus W_r} K_\delta(x-y) dx dy = \int_{W_r} \int_{\mathbb{R}^2 \setminus \mathcal{C}_r} K_\delta(x-y) dx dy + \int_{W_r} \int_{\mathcal{C}} K_\delta(x-y) dx dy.$$

Substituting this identity into (5.2), we find that

$$(5.3) \quad \begin{aligned} \text{Per}_\delta(\mathcal{C}_r, B_R) - \text{Per}_\delta(\mathcal{C}, B_R) &= \text{Per}_\delta(W_r) - 2 \int_{W_r} \int_{\mathcal{C}} K_\delta(x-y) dx dy \\ &= \int_{W_r} \int_{\mathbb{R}^2 \setminus \mathcal{C}_r} K_\delta(x-y) dx dy - \int_{W_r} \int_{\mathcal{C}} K_\delta(x-y) dx dy. \end{aligned}$$

Now, given $x = (x_1, x_2) \in W_r$, we have that $x \in \partial \mathcal{C}_{r(x)}$, with $r(x) := |x_2| \in (0, r]$, where the notation of (1.13) has been used. Then, by Lemma 3.1, we have that

$$(5.4) \quad H_{\mathcal{C}_{r(x)}}^\delta(x) \leq -2\Psi_\delta(r(x)),$$

where Ψ_δ is as in (1.12) with K_δ in place of K , that is

$$\Psi_\delta(s) := \int_{B_{s/4}(7s/4, 0)} K_\delta(x) dx \geq 0.$$

We write (5.4) as

$$\begin{aligned} -2\Psi_\delta(r(x)) &\geq \int_{\mathbb{R}^2 \setminus \mathcal{C}_{r(x)}} K_\delta(x-y) dy - \int_{\mathcal{C}_{r(x)}} K_\delta(x-y) dy \\ &= \int_{\mathbb{R}^2 \setminus \mathcal{C}_r} K_\delta(x-y) dy - \int_{\mathcal{C}} K_\delta(x-y) dy - \int_{W_{r(x)}} K_\delta(x-y) dy \\ &= \int_{\mathbb{R}^2 \setminus \mathcal{C}_r} K_\delta(x-y) dy + \int_{W_r \setminus W_{r(x)}} K_\delta(x-y) dy - \int_{\mathcal{C}} K_\delta(x-y) dy - \int_{W_{r(x)}} K_\delta(x-y) dy. \end{aligned}$$

Therefore, integrating over $x \in W_r$,

$$(5.5) \quad \begin{aligned} &\int_{W_r} \int_{\mathbb{R}^2 \setminus \mathcal{C}_r} K_\delta(x-y) dx dy - \int_{W_r} \int_{\mathcal{C}} K_\delta(x-y) dx dy \\ &\leq \int_{W_r} \int_{W_{r(x)}} K_\delta(x-y) dx dy - \int_{W_r} \int_{W_r \setminus W_{r(x)}} K_\delta(x-y) dx dy - 2 \int_{W_r} \Psi_\delta(r(x)) dx \\ &= 2 \int_{W_r} \int_{W_{r(x)}} K_\delta(x-y) dx dy - \int_{W_r} \int_{W_r} K_\delta(x-y) dx dy - 2 \int_{W_r} \Psi_\delta(r(x)) dx. \end{aligned}$$

We now observe that

$$W_r = \{x \in \mathbb{R}^2 \text{ s.t. } |x_2| > |x_1| \text{ and } |x_2| < r\},$$

and thus

$$\begin{aligned}
& 2 \int_{W_r} \int_{W_{r(x)}} K_\delta(x-y) dx dy \\
&= \int_{x \in W_r} \left(\int_{y \in W_{r(x)}} K_\delta(x-y) dy \right) dx + \int_{y \in W_r} \left(\int_{x \in W_{r(y)}} K_\delta(x-y) dx \right) dy \\
&= \int_{\{|x_1| < |x_2| < r\}} \left(\int_{\{|y_1| < |y_2| < r(x)\}} K_\delta(x-y) dy \right) dx + \int_{\{|y_1| < |y_2| < r\}} \left(\int_{\{|x_1| < |x_2| < r(y)\}} K_\delta(x-y) dx \right) dy \\
&= \int_{\{|x_1| < |x_2| < r\}} \left(\int_{\{|y_1| < |y_2| < |x_2|\}} K_\delta(x-y) dy \right) dx + \int_{\{|y_1| < |y_2| < r\}} \left(\int_{\{|x_1| < |x_2| < |y_2|\}} K_\delta(x-y) dx \right) dy \\
&= \int_{\{|x_1| < |x_2| < r\}} \left(\int_{\{|y_1| < |y_2| < |x_2|\}} K_\delta(x-y) dy \right) dx + \int_{\{|x_1| < |x_2| < r\}} \left(\int_{\{\max\{|y_1|, |x_2|\} < |y_2| < r\}} K_\delta(x-y) dy \right) dx \\
&= \int_{\{|x_1| < |x_2| < r\}} \left(\int_{|y_1| < |y_2| < r} K_\delta(x-y) dx \right) dy.
\end{aligned}$$

Hence, plugging this information into (5.5), we conclude that

$$\int_{W_r} \int_{\mathbb{R}^2 \setminus \mathcal{C}_r} K_\delta(x-y) dx dy - \int_{W_r} \int_{\mathcal{C}} K_\delta(x-y) dx dy \leq -2 \int_{W_r} \Psi_\delta(r(x)) dx.$$

This and (5.3) give that

$$(5.6) \quad \text{Per}_\delta(\mathcal{C}_r, B_R) - \text{Per}_\delta(\mathcal{C}, B_R) \leq -2 \int_{W_r} \Psi_\delta(r(x)) dx.$$

Now, as $\delta \searrow 0$, we have that $\text{Per}_\delta(\mathcal{C}_r, B_R) \rightarrow \text{Per}_K(\mathcal{C}_r, B_R)$ and $\text{Per}_\delta(\mathcal{C}, B_R) \rightarrow \text{Per}_K(\mathcal{C}, B_R)$, by Dominated Convergence Theorem, see [8]. Moreover, $\Psi_\delta(s) \rightarrow \Psi(s) = \int_{B_{s/4}(7s/4, 0)} K(x) dx$ a.e. and in $L^1(0, 1)$ by Dominated Convergence Theorem (observe that $\Psi \in L^1(0, 1)$ by assumption (1.3)).

So, letting $\delta \searrow 0$ in (5.6), we end up with

$$(5.7) \quad \text{Per}_K(\mathcal{C}_r, B_R) - \text{Per}_K(\mathcal{C}, B_R) \leq -2 \int_{W_r} \Psi(|x_2|) dx.$$

Recalling that K is not identically zero, we take a Lebesgue point $\tau_0 \in (0, +\infty)$ such that $K_0(\tau_0) > 0$. Then,

$$\lim_{\varepsilon \searrow 0} \frac{1}{2\varepsilon} \int_{\tau_0 - \varepsilon}^{\tau_0 + \varepsilon} K_0(\tau) d\tau = K_0(\tau_0) > 0.$$

Consequently, we take $\varepsilon_0 > 0$ such that for all $\varepsilon \in (0, \varepsilon_0]$ we have that

$$(5.8) \quad \int_{\tau_0 - \varepsilon}^{\tau_0 + \varepsilon} K_0(\tau) d\tau \geq \varepsilon K_0(\tau_0).$$

Then, if $\bar{\varepsilon} := \min\{\varepsilon_0, \frac{\tau_0}{100}\}$ and $r \in [\frac{4\tau_0}{7} - \frac{\bar{\varepsilon}}{14}, \frac{4\tau_0}{7} + \frac{\bar{\varepsilon}}{14}]$, we have that

$$(5.9) \quad \begin{aligned} \frac{7r}{4} + \frac{r}{8} &= \frac{15r}{8} \geq \frac{15\tau_0}{8} - \frac{15\bar{\varepsilon}}{8} \geq \tau_0 + \bar{\varepsilon} \\ \text{and } \frac{7r}{4} - \frac{r}{8} &= \frac{13r}{8} \leq \frac{13\tau_0}{8} + \frac{13\bar{\varepsilon}}{8} \leq \tau_0 - \bar{\varepsilon}. \end{aligned}$$

Now we cover the ring $A_r := B_{(7r/4)+(r/8)} \setminus B_{(7r/4)-(r/8)}$ by N_0 balls of radius $r/4$ centered at $\partial B_{7r/4}$, with N_0 independent of r . Then

$$\begin{aligned}
& \frac{13\pi r}{4} \int_{(7r/4)-(r/8)}^{(7r/4)+(r/8)} K_0(\tau) d\tau \leq 2\pi \int_{(7r/4)-(r/8)}^{(7r/4)+(r/8)} \tau K_0(\tau) d\tau \\
&= \int_{A_r} K_0(|x|) dx \leq N_0 \int_{B_{r/4}(7r/4, 0)} K_0(|x|) dx = N_0 \Psi(r),
\end{aligned}$$

thanks to (1.12).

Using this, (5.8) and (5.9), we obtain that, for any $r \in [\frac{4\tau_0}{7} - \frac{\bar{\varepsilon}}{14}, \frac{4\tau_0}{7} + \frac{\bar{\varepsilon}}{14}]$,

$$\begin{aligned}
(5.10) \quad \Psi(r) &\geq \frac{13\pi r}{4N_0} \int_{(7r/4)-(r/8)}^{(7r/4)+(r/8)} K_0(\tau) d\tau \\
&\geq \frac{\tau_0}{4N_0} \int_{\tau_0-\bar{\varepsilon}}^{\tau_0+\bar{\varepsilon}} K_0(\tau) d\tau \\
&\geq \frac{\bar{\varepsilon}\tau_0 K_0(\tau_0)}{4N_0} \\
&=: \bar{c}.
\end{aligned}$$

Then, if $r_0 := \frac{4\tau_0}{7} + \frac{\bar{\varepsilon}}{14}$, we have that

$$W_{r_0} \supset \left(0, \frac{4\tau_0}{7} - \frac{\bar{\varepsilon}}{14}\right) \times \left(\frac{4\tau_0}{7} - \frac{\bar{\varepsilon}}{14}, \frac{4\tau_0}{7} + \frac{\bar{\varepsilon}}{14}\right)$$

and therefore

$$\int_{W_{r_0}} \Psi(|x_2|) dx \geq \left(\frac{4\tau_0}{7} - \frac{\bar{\varepsilon}}{14}\right) \int_{\frac{4\tau_0}{7}-\frac{\bar{\varepsilon}}{14}}^{\frac{4\tau_0}{7}+\frac{\bar{\varepsilon}}{14}} \Psi(x_2) dx_2 \geq \frac{\bar{c}\bar{\varepsilon}}{7} \left(\frac{4\tau_0}{7} - \frac{\bar{\varepsilon}}{14}\right),$$

where (5.10) has been used in the last inequality. In particular,

$$\int_{W_{r_0}} \Psi(|x_2|) dx > 0,$$

which combined with (5.7) implies that claim in (5.1) with $r := r_0$.

Then, in light of (5.1), we get that \mathcal{C} is not a K -minimal set, thus completing the proof of Proposition 1.9. \square

6. STRICTLY STARSHAPED DOMAINS AND PROOF OF PROPOSITION 1.10

Proof of Proposition 1.10. We observe that, due to assumption in (1.28), for every $\lambda > 0$, we have that there exists $\delta_\lambda > 0$ such that the distance between ∂E and $\partial(\lambda E)$ is at least δ_λ . Therefore, for any $\lambda > 1$, from Corollary A.8 and Lemma A.13, we deduce that

$$E^+(\lambda^{1+s}t) \subseteq E_\lambda^-(\lambda^{1+s}t) = \lambda E^-(t).$$

Then for $\lambda > 1$,

$$|\text{int}(E^+(t)) \setminus \overline{E^-(t)}| \leq |\text{int}(E^+(t)) \setminus \lambda^{-1}E^+(\lambda^{1+s}t)| = |\text{int}(E^+(t))| - \lambda^{-1}|E^+(\lambda^{1+s}t)|.$$

Also, by Proposition A.12,

$$\liminf_{\lambda \searrow 1} |E^+(\lambda^{1+s}t)| \geq |\text{int}(E^+(t))|.$$

Therefore we get

$$|\text{int}(E^+(t)) \setminus \overline{E^-(t)}| \leq \limsup_{\lambda \searrow 1} |\text{int}(E^+(t))| - \lambda^{-1}|E^+(\lambda^{1+s}t)| = |\text{int}(E^+(t))| - \liminf_{\lambda \searrow 1} \lambda^{-1}|E^+(\lambda^{1+s}t)| \leq 0.$$

This gives the desired statement. \square

7. PERTURBED DOUBLE DROPLET AND PROOF OF THEOREM 1.11

Recalling the notation in (1.29), given $r \in (0, \frac{1}{2})$ we set


$$(7.1) \quad \mathcal{G}_r := [-r, r]^2 \cup \mathcal{G}_0,$$

where \mathcal{G}_0 is the union of \mathcal{B}^+ , which is the convex envelope between $B_1(\sqrt{2}, 0)$ and the origin, and \mathcal{B}^- , which is the convex envelope between $B_1(-\sqrt{2}, 0)$ and the origin, see Figure 12.

Now, fixed $\delta \in (0, r)$, we denote by \mathcal{B}_δ^+ the convex envelope between $B_{1-\delta}(\sqrt{2}, 0)$ and the origin, and \mathcal{B}_δ^- the convex envelope between $B_{1-\delta}(-\sqrt{2}, 0)$ and the origin. We let

$$\mathcal{G}_{\delta,r} := ([-2r, 2r] \times [-r, r]) \cup \mathcal{B}_\delta^+ \cup \mathcal{B}_\delta^-.$$

Then we can estimate the K -curvature of $\mathcal{G}_{\delta,r}$ as follows:

FIGURE 12. The set \mathcal{G}_r .

Lemma 7.1. *Assume that (1.2), (1.3) and (1.27) hold true. Then, there exists $c_{\sharp} \in (0, 1)$ such that the following statement holds true. If $r \in (0, c_{\sharp})$ and $\delta \in (0, c_{\sharp}^4 r)$, then*

$$(7.2) \quad H_{\mathcal{G}_{\delta,r}}^s(p) \leq \frac{1}{c_{\sharp}}$$

for any $p \in \partial\mathcal{G}_{\delta,r}$. In addition, for any $p \in (\partial\mathcal{G}_{\delta,r}) \cap ([-2r, 2r] \times [-r, r])$,

$$(7.3) \quad H_{\mathcal{G}_{\delta,r}}^s(p) \leq -\frac{c_{\sharp}}{r^s}.$$

Proof. Let $\alpha(\delta)$ the angle at $x = 0$ in \mathcal{B}_{δ}^+ . Observe that when $\delta = 0$, this angle is $\pi/2$ and moreover there exist δ_0 and $C_0 > 0$ such that $|\alpha(\delta) - \frac{\pi}{2}| \leq C_0\delta$, for all $0 < \delta < \delta_0$. In particular we may assume that $\alpha(\delta) \geq \pi/3$. We fix then $\delta \leq r < \delta_0$.

First of all note that for all $p = (p_1, p_2) \in \partial\mathcal{G}_{\delta,r}$, with $p_1 \geq \sqrt{2} - \frac{(1-\delta)^2}{\sqrt{2}}$ (resp. $p_1 \leq -\sqrt{2} + \frac{(1-\delta)^2}{\sqrt{2}}$), then $p \in \partial B_{1-\delta}(\sqrt{2}, 0)$ (resp. $p \in \partial B_{1-\delta}(-\sqrt{2}, 0)$), and then

$$H_{\mathcal{G}_{\delta,r}}^s(p) \leq H_{B_{1-\delta}(\sqrt{2}, 0)}^s(p) = c(1)(1-\delta)^{-s} \quad \left(\text{resp. } H_{\mathcal{G}_{\delta,r}}^s(p) \leq H_{B_{1-\delta}(-\sqrt{2}, 0)}^s(p) = c(1)(1-\delta)^{-s}\right)$$

where $c(1) = H_{B_1}^s$.

We take $c_{\sharp} \in (0, 1)$ to be taken conveniently small in what follows. We notice that $\mathcal{S} := (\partial\mathcal{G}_{\delta,r}) \cap \{|x_2| = r\}$ consists of four points. We take $p = (p_1, p_2) \in \partial\mathcal{G}_{\delta,r}$ such that there exists $q \in \mathcal{S}$ such that $|p - q| < c_{\sharp}r$ (see e.g. Figure 13 for a possible configuration).

Then,

$$(7.4) \quad \begin{aligned} & \lim_{\varepsilon \searrow 0} \int_{B_{\sqrt{c_{\sharp}}r}(p) \setminus B_{\varepsilon}(p)} \left(\chi_{\mathbb{R}^2 \setminus \mathcal{G}_{\delta,r}}(y) - \chi_{\mathcal{G}_{\delta,r}}(y) \right) \frac{1}{|p - y|^{2+s}} dy \\ & \leq - \iint_{(0, \pi/6) \times (c_{\sharp}r, \sqrt{c_{\sharp}}r)} \frac{1}{\varrho^{1+s}} d\vartheta d\rho = -\frac{\pi}{6s} \frac{1}{c_{\sharp}^{s/2} r^s} \left(\frac{1}{c_{\sharp}^{s/2}} - 1 \right), \end{aligned}$$

while

$$\int_{\mathbb{R}^2 \setminus B_{\sqrt{c_{\sharp}}r}(p)} \left(\chi_{\mathbb{R}^2 \setminus \mathcal{G}_{\delta,r}}(y) - \chi_{\mathcal{G}_{\delta,r}}(y) \right) \frac{1}{|p - y|^{2+s}} \leq 2\pi \int_{\sqrt{c_{\sharp}}r}^{+\infty} \frac{1}{\rho^{1+s}} d\rho = \frac{2\pi}{s} \frac{1}{c_{\sharp}^{s/2} r^s}.$$

As a consequence,

$$H_{\mathcal{G}_{\delta,r}}^K(p) \leq -\frac{\pi}{6s} \frac{1}{c_{\sharp}^{s/2} r^s} \left(\frac{1}{c_{\sharp}^{s/2}} - 1 \right) + \frac{2\pi}{s} \frac{1}{c_{\sharp}^{s/2} r^s} \leq -c_{\sharp} \frac{1}{r^s}$$

as long as c_{\sharp} is sufficiently small, which implies (7.3) (and also (7.2)) in this case.

Now consider $p \in \partial\mathcal{G}_{\delta,r}$ such that $p_2 \neq r$ and $d(p, \mathcal{S}) \geq c_{\sharp}r$. If $p \in \partial B_{1-\delta}(\pm\sqrt{2}, 0)$ we are ok, and in the other case, note that we can define a set \mathcal{G}' with $C^{1,1}$ -boundary (uniformly in δ and r) such that $\mathcal{G}_{\delta,r} \subset \mathcal{G}'$ and $\mathcal{G}' \setminus B_{1/8} = \mathcal{G}_{\delta,r} \setminus B_{1/8}$. Then, we obtain that

$$C' \geq H_{\mathcal{G}'}^K(p) \geq H_{\mathcal{G}_{\delta,r}}^K(p) - C'',$$

for some $C', C'' > 0$, depending only on the local $C^{1,1}$ -norms of the boundary of \mathcal{G}' , and this gives (7.2) in this case.




FIGURE 13. .

Finally note that $\mathcal{G}_{\delta,r} \subseteq \mathcal{C}_r$, where \mathcal{C}_r is the perturbed cross is defined in (1.13). So, if $p \in \partial\mathcal{G}_{\delta,r} \cap ([-r, r] \times [-r, r])$, then $p \in \partial\mathcal{C}_r$. Moreover by Lemma 3.1 and the definition of Ψ in (1.12)

$$H_{\mathcal{C}_r}^s(p) \leq -2\Psi(r) = -C \frac{1}{r^s}$$

where $C > 0$ is a universal constant. In this case, we notice that $\mathcal{G}_{\delta,r}$ and \mathcal{C}_r coincide in B_r , and, outside such a neighborhood of the origin, they differ by four portions of cones (passing in the vicinity of \mathcal{S}) with opening bounded by $C_0\delta$. That is, if we set

$$\mathcal{D}_{\delta,r} := (\mathcal{G}_{\delta,r} \setminus \mathcal{C}_r) \cup (\mathcal{C}_r \setminus \mathcal{G}_{\delta,r}),$$

we have that

$$\int_{\mathcal{D}_{\delta,r}} \frac{dy}{|p-y|^{2+s}} \leq C_2 \left[\iint_{(0,C_1\delta) \times (c_{\sharp}r/2, 10r]} \frac{\rho d\vartheta d\rho}{(c_{\sharp}r/2)^{2+s}} + \iint_{(0,C_1\delta) \times (10r, +\infty)} \frac{\rho d\vartheta d\rho}{\rho^{2+s}} \right] \leq \frac{C_3 \delta}{c_{\sharp}^{2+s} r^s} \leq \frac{c_{\sharp}}{r^s},$$

thanks to our assumption on δ . Consequently

$$|H_{\mathcal{G}_{\delta,r}}^K(p) - H_{\mathcal{C}_r}^K(p)| \leq \frac{c_{\sharp}}{r^s}$$

and so, making use of (1.15) and (1.33),

$$H_{\mathcal{G}_{\delta,r}}^K(p) \leq H_{\mathcal{C}_r}^K(p) + \frac{c_{\sharp}}{r^s} \leq -\frac{c_*}{r^s} + \frac{c_{\sharp}}{r^s} \leq -\frac{c_*}{2r^s},$$

for a suitable $c_* > 0$, as long as $c_{\sharp} > 0$ is sufficiently small. This establishes (7.3) (and also (7.2)) in this case. \square

With these auxiliary computations, we can now complete the proof of Theorem 1.11, by arguing as follows.

Proof of Theorem 1.11. Let $c_{\sharp} > 0$ be as in Lemma 7.1, $0 < \varepsilon < c_{\sharp}/2$ and $c_* := ((c_{\sharp} - \varepsilon)(1+s))^{1/(1+s)}$. We define $r(t)$ such that $\dot{r}(t) = (c_{\sharp} - \varepsilon)r(t)^{-s}$, with $r(0) = 0$. So, we have that $r(t) = c_* t^{1/(1+s)}$. Let also

$$\delta(t) := \frac{1}{c_{\sharp}\varepsilon} \int_0^t \frac{d\tau}{r(\tau)} = \frac{1+s}{(c_{\sharp} - \varepsilon)c_* s} t^{s/(1+s)}.$$

We now estimate the outer normal velocity of $\mathcal{G}_{\delta(t), r(t)}$ via Lemma 7.1. First of all, from (7.3) at $p \in (\partial\mathcal{G}_{\delta(t), r(t)}) \cap \{|x_2| = r(t), |x_1| < \sqrt{2}\}$ we get

$$\dot{r}(t) = \frac{c_{\sharp} - \varepsilon}{(r(t))^s} \leq -H_{\mathcal{G}_{\delta(t), r(t)}}^s(p) - \frac{\varepsilon}{c_*^s t^{s/(1+s)}}.$$

Moreover, the shrinking velocity at $x \in (\partial\mathcal{G}_{\delta(t),r(t)}) \setminus \{|x_2| = r(t)\}$ is at least $r(t)\dot{\delta}(t) = 1/(c_{\sharp} - \varepsilon)$. This implies that at every $x \in (\partial\mathcal{G}_{\delta(t),r(t)}) \setminus \{|x_2| = r(t)\}$ we get

$$\partial_t x \cdot \nu(x) \leq -\frac{1}{c_{\sharp} - \varepsilon} \leq -H_{\mathcal{G}_{\delta(t),r(t)}}^s(x) - \frac{\varepsilon}{c_{\sharp}(c_{\sharp} - \varepsilon)}.$$

by (7.2). Therefore, by Proposition A.10, we get that

$$(7.5) \quad B_{c_{\star} t^{1/(1+s)}} \subseteq \mathcal{G}_{\delta(t),r(t)} \subseteq \mathcal{G}^+(t).$$

Conversely, since \mathcal{G} is contained in the cross \mathcal{C} , it follows from Corollary A.8 and Theorem 1.5 that

$$B_{c_o t^{1/(1+s)}} \subseteq (\mathbb{R}^2 \setminus \mathcal{C})^+(t) \subseteq (\mathbb{R}^2 \setminus \mathcal{G})^+(t).$$

From this and (7.5) it follows that

$$B_{\hat{c} t^{1/(1+s)}} \subseteq \mathcal{G}^+(t) \cap (\mathbb{R}^2 \setminus \mathcal{G})^+(t) = \Sigma_{\mathcal{G}}(t)$$

with $\hat{c} := \min\{c_{\star}, c_o\}$, which proves (1.31). \square

8. PERTURBATION OF TANGENT BALLS AND PROOF OF THEOREM 1.13

The idea to prove Theorem 1.13 is to construct inner barriers using ‘‘almost tangent’’ balls and take advantage of the scale invariance given by the homogeneous kernels in (1.27). For this, given $\delta \in [0, \frac{1}{8}]$, we consider the set

$$\mathcal{Z}_{\delta,r} := B_r((1+\delta)r, 0) \cup B_r((-1-\delta)r, 0).$$

Then, we have that the nonlocal curvature of $\mathcal{Z}_{\delta,r}$ is always controlled from above by that of the ball, and it becomes negative in the vicinity of the origin. More precisely:

Lemma 8.1. *Assume (1.27). Then, for any $p \in \partial\mathcal{Z}_{\delta,r}$ we have that*

$$(8.1) \quad H_{\mathcal{Z}_{\delta,r}}^K(p) \leq \frac{C}{r^s},$$

for some $C > 0$. In addition, there exists $c \in (0, 1)$ such that if $\delta \in (0, c^2)$ and $p \in (\partial\mathcal{Z}_{\delta,r}) \cap B_{cr}$ then

$$(8.2) \quad H_{\mathcal{Z}_{\delta,r}}^K(p) \leq -c.$$

Proof. Notice that $\partial\mathcal{Z}_{\delta,r} \subseteq (\partial B_r((1+\delta)r, 0)) \cup (\partial B_r((-1-\delta)r, 0))$. Moreover, $\mathcal{Z}_{\delta,r} \supseteq B_r((1+\delta)r, 0)$, as well as $\mathcal{Z}_{\delta,r} \supseteq B_r((-1-\delta)r, 0)$, hence, in view of (1.4), the nonlocal curvature of $\mathcal{Z}_{\delta,r}$ is less than or equal to that of B_r , which proves (8.1).

Now we prove (8.2). For this, up to scaling, we assume that $r := 1$ and we take $p \in (\partial\mathcal{Z}_{\delta,1}) \cap B_c$. Without loss of generality, we also suppose that $p_1, p_2 > 0$ and we observe that

$$(8.3) \quad B_c(-2c, 0) \subseteq B_1((-1-\delta), 0),$$

as long as c is small enough. Indeed if $x \in B_c(2c, 0)$ then we can write $x = -2ce_1 + ce$, for some $e \in \mathbb{S}^1$, and so

$$|x - (-1-\delta)e_1| = |(1+\delta-2c)e_1 + ce| \leq |1+\delta-2c| + c = (1+\delta-2c) + c = 1+\delta-c \leq 1+c^2-c < 1.$$

This proves (8.3).

Hence, from (1.4) and (8.3), the nonlocal curvature of $\mathcal{Z}_{\delta,1}$ at p is less than or equal to the nonlocal curvature of $B_r((1+\delta), 0)$, which is bounded by some $C > 0$, minus the contribution coming from $B_c(-2c, 0)$. That is,

$$(8.4) \quad -H_{\mathcal{Z}_{\delta,r}}^K(p) \geq -C + \int_{B_c(-2c,0)} \frac{dx}{|x-p|^{2+s}} = -C + \int_{B_c(2c+p_1,p_2)} \frac{dx}{|y|^{2+s}}.$$

Also, if $y \in B_c(2c+p_1,p_2)$, we have that $|y| \leq |y-2ce_1-p| + |2ce_1+p| \leq c+2c+|p| \leq 4c$, and so

$$\int_{B_c(2c+p_1,p_2)} \frac{dx}{|y|^{2+s}} \geq \frac{c_0 c^2}{c^{2+s}} = \frac{c_0}{c^s},$$

for some $c_0 > 0$. So we insert this information into (8.4) and we obtain

$$-H_{\mathcal{Z}_{\delta,r}}^K(p) \geq -C + \frac{c_0}{c^s} \geq \frac{c_0}{2c^s}$$

as long as c is sufficiently small. This completes the proof of (8.2), as desired. \square

From Lemma 8.1, we can control the geometric flow of the double tangent balls from inside with barriers that shrink the sides of the picture and make the origin emanate some mass:

Lemma 8.2. *There exist $\delta_0 \in (0, 1)$, and $\bar{C} > 0$ such that if $\delta \in (0, \delta_0)$, then*

$$\mathcal{O}^-(\delta) \supseteq \bigcup_{\sigma \in (-\delta^2, \delta^2)} (B_{1-\bar{C}\delta}(1 - \bar{C}\delta, 0) \cup B_{1-\bar{C}\delta}(-1 + \bar{C}\delta, 0) + \sigma e_2).$$

Proof. Fix $\varepsilon \in (0, 1)$, to be taken arbitrarily small in what follows. Let $\mu \in [0, \sqrt{\varepsilon}]$ and let, for any $t \in [0, (1 - \varepsilon)/C_0]$,

$$\varepsilon_\mu(t) := \varepsilon - \mu t \quad \text{and} \quad r(t) := 1 - \varepsilon - C_0 t,$$

with $C_0 > 0$ to be chosen conveniently large. We consider an inner barrier consisting in two balls of radius $r(t)$ which, for any $t \in [0, (1 - \varepsilon)/C_0]$, remain at distance $2\varepsilon(t)$. Namely, we set

$$(8.5) \quad \mathcal{F}_{\varepsilon, \mu}(t) := B_{r(t)}(r(t) + \varepsilon_\mu(t), 0) \cup B_{r(t)}(-r(t) - \varepsilon_\mu(t), 0).$$

Notice that

$$(8.6) \quad \mathcal{O} \supseteq \mathcal{F}_{\varepsilon, \mu}(0) + \sigma e_2 \quad \text{for any } \sigma \in (-\varepsilon, \varepsilon) \quad d(\mathcal{O}, \mathcal{F}_{\varepsilon, \mu}(0) + \sigma e_2) > 0.$$

We also observe that the vectorial velocity of this set is the superposition of a normal velocity $-\dot{r}\nu$, being ν the interior normal, and a translation velocity $\pm(\dot{r} + \dot{\varepsilon}_\mu)e_1$, with the plus sign for the ball on the right and the minus sign for the ball on the left. The normal velocity of this set is therefore equal to

$$(8.7) \quad \left(-\dot{r}\nu \pm (\dot{r} + \dot{\varepsilon}_\mu)e_1 \right) \cdot \nu = -\dot{r} \pm (\dot{r} + \dot{\varepsilon}_\mu)\nu_1 = C_0(1 \mp \nu_1) \mp \mu.$$

Now, taken a point p on $\partial\mathcal{F}_{\varepsilon, \mu}(t)$, we distinguish two cases. Either $p \in B_c$, where c is the one given in Lemma 8.1, or $p \in \mathbb{R}^2 \setminus B_c$. In the first case, we have that

$$C_0(1 \mp \nu_1) \mp \mu \geq C_0(1 - |\nu_1|) - \mu \geq 0 - \mu \geq -\sqrt{\varepsilon} > -c,$$

This and (8.7) give that the normal velocity of $\mathcal{F}_{\varepsilon, \mu}(t)$ at p is larger than $-c$, and therefore greater than $H_{z_{\delta, r}}^K(p)$, thanks to (8.2).

If instead $p \in \mathbb{R}^2 \setminus B_c$, we have that $|\nu_1(p)| \leq 1 - c_0$, for a suitable $c_0 \in (0, 1)$, depending on c , and therefore

$$C_0(1 \mp \nu_1) \mp \mu \geq C_0(1 - |\nu_1|) - \mu \geq C_0 c_0 - \mu \geq C_0 c_0 - 1 \geq \frac{C_0 c_0}{2} \geq \frac{C_0 c_0}{2^{s+1} (r(t))^s},$$

as long as C_0 is sufficiently large. This and (8.7) give that the inner normal velocity of $\mathcal{F}_{\varepsilon, \mu}(t)$ at p is strictly larger than $\frac{C_0 c_0}{2^{s+1} (r(t))^s}$, which, if C_0 is chosen conveniently big, is in turn strictly larger than $H_{z_{\delta, r}}^K(p)$, thanks to (8.1).

In any case, we have shown that the inner normal velocity of $\mathcal{F}_{\varepsilon, \mu}(t)$ at p is strictly larger than $H_{z_{\delta, r}}^K(p)$. This implies that $\mathcal{F}_{\varepsilon, \mu}(t)$ is a strict subsolution according to Proposition A.10.

Then, by (8.6) and Proposition A.10

$$(8.8) \quad \mathcal{O}^-(t) \supseteq \bigcup_{\sigma \in (-\varepsilon, \varepsilon)} (\mathcal{F}_{\varepsilon, \mu}(t) + \sigma e_2),$$

for any $t \in [0, (1 - \varepsilon)/C_0]$.

Now, taking $\mu := \sqrt{\varepsilon}$ in (8.5), we see that

$$\mathcal{F}_{\varepsilon, \sqrt{\varepsilon}}(t) = B_{1-\varepsilon-C_0 t}(1 - \sqrt{\varepsilon}t - C_0 t, 0) \cup B_{1-\varepsilon-C_0 t}(-1 + \sqrt{\varepsilon}t - C_0 t, 0)$$

for all $t \in [0, (1 - \varepsilon)/C_0]$. In particular, taking $t := \sqrt{\varepsilon}$,

$$\mathcal{F}_{\varepsilon, \sqrt{\varepsilon}}(\sqrt{\varepsilon}) = B_{1-\varepsilon-C_0\sqrt{\varepsilon}}(1 - \varepsilon - C_0\sqrt{\varepsilon}, 0) \cup B_{1-\varepsilon-C_0\sqrt{\varepsilon}}(-1 + \varepsilon - C_0\sqrt{\varepsilon}, 0),$$

and the latter are two tangent balls at the origin. From this and (8.8), we deduce that

$$\mathcal{O}^-(\sqrt{\varepsilon}) \supseteq \bigcup_{\sigma \in (-\varepsilon, \varepsilon)} (B_{1-\varepsilon-C_0\sqrt{\varepsilon}}(1 - \varepsilon - C_0\sqrt{\varepsilon}, 0) \cup B_{1-\varepsilon-C_0\sqrt{\varepsilon}}(-1 + \varepsilon - C_0\sqrt{\varepsilon}, 0) + \sigma e_2),$$

and this implies the desired result by choosing $\delta := \sqrt{\varepsilon}$ and $\bar{C} := 2(C_0 + 1)$. \square

We can now complete the proof of Theorem 1.13 in the following way:

Proof of Theorem 1.13. We observe that, in the setting of Lemma A.13, the result in Lemma 8.2 can be written as

$$\mathcal{O}^-(\delta) \supseteq (1 - \bar{C}\delta)\mathcal{O}^+(0) \quad \text{with} \quad d(\mathcal{O}^-(\delta), (1 - \bar{C}\delta)\mathcal{O}^+(0)) \geq \delta^2$$

for all $\delta \in (0, \delta_0)$.

Fix now $C \geq \bar{C}$ and let $\mathcal{U} := (1 - C\delta)\mathcal{O}^+(0)$. Then, by Corollary A.8, we have

$$(8.9) \quad \mathcal{O}^-(t + \delta) \supseteq \mathcal{U}(t)$$

for all $t \geq 0$.

Now, in view of Lemma A.13,

$$\mathcal{U}(t) = (1 - C\delta) \mathcal{O}^+ \left(\frac{t}{(1 - C\delta)^{1+s}} \right)$$

and so, combining with (8.9),

$$\mathcal{O}^-(t + \delta) \supseteq (1 - C\delta) \mathcal{O}^+ \left(\frac{t}{(1 - C\delta)^{1+s}} \right).$$

Consequently, for any $t \geq \delta$, we can estimate the measure of the fattening set as

$$(8.10) \quad \begin{aligned} |\text{int}(\mathcal{O}^+(t)) \setminus \overline{\mathcal{O}^-(t)}| &\leq \left| \text{int}(\mathcal{O}^+(t)) \setminus (1 - C\delta) \mathcal{O}^+ \left(\frac{t - \delta}{(1 - C\delta)^{1+s}} \right) \right| \\ &= |\text{int}(\mathcal{O}^+(t))| - \left| (1 - C\delta) \mathcal{O}^+ \left(\frac{t - \delta}{(1 - C\delta)^{1+s}} \right) \right| \\ &= |\text{int}(\mathcal{O}^+(t))| - (1 - C\delta)^2 \left| \mathcal{O}^+ \left(\frac{t - \delta}{(1 - C\delta)^{1+s}} \right) \right|. \end{aligned}$$

We now fix $t_0 \geq \delta$ and choose $C = C(t_0) \geq \bar{C}$ such that

$$t \leq \frac{t - \delta}{(1 - C\delta)^{1+s}} \quad \text{for all } t \geq t_0.$$

So, by Proposition A.12, we get that

$$\liminf_{\delta \searrow 0} \left| \mathcal{O}^+ \left(\frac{t - \delta}{(1 - C\delta)^{1+s}} \right) \right| \geq |\text{int}(\mathcal{O}^+(t))|.$$

This and (8.10) yield that, for $t \geq t_0$,

$$\begin{aligned} |\text{int}(\mathcal{O}^+(t)) - \overline{\mathcal{O}^-(t)}| &\leq \limsup_{\delta \searrow 0} |\text{int}(\mathcal{O}^+(t))| - (1 - C\delta)^2 \left| \mathcal{O}^+ \left(\frac{t - \delta}{(1 - C\delta)^{1+s}} \right) \right| \\ &= |\text{int}(\mathcal{O}^+(t))| - \liminf_{\delta \searrow 0} (1 - C\delta)^2 \left| \mathcal{O}^+ \left(\frac{t - \delta}{(1 - C\delta)^{1+s}} \right) \right| \leq 0. \end{aligned}$$

Since t_0 was chosen arbitrarily, this completes the proof of Theorem 1.13. \square

APPENDIX A. VISCOSITY SOLUTIONS AND GEOMETRIC BARRIERS

In this appendix, we recall the existence and uniqueness results about the level set flow associated to the nonlocal evolution (1.1), and we provide some auxiliary results which will be useful in the proof of the main theorems.

Before introducing the level set equation and the notion of viscosity solutions, we briefly discuss the evolution of balls according to the setting in (1.1)–(1.4).

Lemma A.1. *Assume that (1.2) and (1.3) hold true. Then for every $R > 0$ there exists $c(R) > 0$ such that*

$$H_{B_R}^K(x) = c(R) \quad \text{for all } x \in \partial B_R.$$

Moreover the function

$$R \in (0, +\infty) \rightarrow c(R) \in (0, +\infty)$$

is continuous, nonincreasing and such that

$$\lim_{R \rightarrow +\infty} c(R) = 0.$$

Furthermore, if K is a fractional kernel, that is $K(x) = \frac{1}{|x|^{2+s}}$, then $c(R) = c(1)R^{-s}$.

Proof. We observe that, in virtue of (1.2) and (1.4), and the fact that $K(x) \not\equiv 0$, we have that $H_{B_R}^K(x) > 0$ for $x \in \partial B_R$, it does not depend on x , and finally $H_{B_R}^K(x) \geq H_{B_{R'}}^K(x)$ if $R' > R$. Condition (1.3) assures that $H_{B_R}^K(x)$ is finite for every $R > 0$ and also that

$$\lim_{R \rightarrow +\infty} c(R) = 0,$$

see [8, 16]. For the computation in the case of fractional kernels, see [18]. \square

Remark A.2. Using Lemma A.1, we study the evolution of a ball B_R according to the flow in (1.1). Such evolution is given by a ball $B_{R(t)}$, where

$$(A.1) \quad \dot{R}(t) = -c(R(t))$$

with initial datum $R(0) = R$. We define

$$C(R) := \int_1^R \frac{1}{c(s)} ds.$$

Then $C(R)$ is a monotone increasing function, and the solution $R(t)$ to (A.1) is given implicitly by the formula

$$(A.2) \quad C(R(t)) = C(R) - t \quad \text{for all } t > 0, \text{ s.t. } R(t) > 0.$$

Let also

$$T_R := \sup\{t > 0 \mid R(t) > 0\}.$$

By (A.1) and the monotonicity of $c(\cdot)$, it is easy to check that

$$T_R \leq \frac{R}{c(R)}.$$

Moreover, from (A.1) we have that

$$T_R = \lim_{\varepsilon \searrow 0} \int_\varepsilon^R \frac{1}{c(s)} ds = C(R) - \lim_{\varepsilon \searrow 0} C(\varepsilon).$$

If K is a fractional kernel, that is $K(x) = \frac{1}{|x|^{2+s}}$, then $C(R) = \frac{1}{c(1)(s+1)}(R^{s+1} - 1)$ and $T_R = \frac{R^{s+1}}{c(1)(s+1)}$.

We introduce now the notion of viscosity solutions for the level set equation

$$(A.3) \quad \begin{cases} \partial_t u(x, t) + |Du(x, t)| H_{\{y|u(y,t) \geq u(x,t)\}}^K(x) = 0 & \text{for all } x \in \mathbb{R}^2, \quad t > 0, \\ u(x, 0) = u_0(x) & \text{for all } x \in \mathbb{R}^2. \end{cases}$$

For more details, we refer to [8, 16]. The viscosity theory for the classical mean curvature flow is contained in [10], see also [14] for a comprehensive level set approach for classical geometric flows.

Definition A.3 (Viscosity solutions).

- i) An upper semicontinuous function $u : \mathbb{R}^2 \times (0, T) \rightarrow \mathbb{R}$ is a viscosity subsolution of (A.3) if, for every smooth test function ϕ such that $u - \phi$ admits a global maximum at (x, t) , we have that either $\partial_t \phi(x, t) \leq 0$ if $D\phi(x, t) = 0$, or

$$\partial_t \phi(x, t) + |D\phi(x, t)| H_{\{y|\phi(y,t) \geq \phi(x,t)\}}^K(x) \leq 0$$

if $D\phi(x, t) \neq 0$.

- ii) A lower semicontinuous function $u : \mathbb{R}^2 \times (0, T) \rightarrow \mathbb{R}$ is a viscosity supersolution of (A.3) if, for every smooth test function ϕ such that $u - \phi$ admits a global minimum at (x, t) , we have that either $\partial_t \phi(x, t) \geq 0$ if $D\phi(x, t) = 0$, or

$$\partial_t \phi(x, t) + |D\phi(x, t)| H_{\{y|\phi(y,t) > \phi(x,t)\}}^K(x) \geq 0$$

if $D\phi(x, t) \neq 0$.

- iii) A continuous function $u : \mathbb{R}^2 \times (0, T) \rightarrow \mathbb{R}$ is a solution to (A.3) if it is both a subsolution and a supersolution.

Remark A.4. It is easy to verify that any smooth subsolution (respectively supersolution) is in particular a viscosity subsolution (respectively supersolution).

Now, we recall the Comparison Principle and the existence and uniqueness results for viscosity solutions to (A.3).

Theorem A.5. *Suppose that u_0 is a bounded and uniformly continuous function. Let u (respectively v) be a bounded viscosity subsolution (respectively supersolution) of (A.3). If $u(x, 0) \leq u_0(x) \leq v(x, 0)$ for any $x \in \mathbb{R}^2$, then $u \leq v$ on $\mathbb{R}^2 \times [0, +\infty)$.*

In particular, there exists a unique continuous viscosity solution u to (A.3) such that $u(x, 0) = u_0(x)$ for any $x \in \mathbb{R}^2$.

Moreover if u_0 is Lipschitz continuous then $u(\cdot, t)$ is Lipschitz continuous, uniformly with respect to t , and

$$|u(x, t) - u(y, t)| \leq \|Du_0\|_\infty |x - y|,$$

for all $x, y \in \mathbb{R}^2$ and $t > 0$.

Proof. For the proof of the existence and uniqueness result, and for the Comparison Principle, we refer to [16, Theorems 2 and 3], see also [8].

Finally the Lipschitz continuity is a consequence of the Comparison Principle. Indeed, for any $h \in \mathbb{R}^2$, we define

$$v_\pm(x, t) := u(x + h, t) \pm \|Du_0\|_\infty |h|.$$

Then, if u is a viscosity solution to (A.3), we have that also v_+ and v_- are viscosity solutions to the same equation. Moreover,

$$v_-(x, 0) = u_0(x + h) - \|Du_0\|_\infty |h| \leq u_0(x) = u(x, 0) \leq u_0(x + h) + \|Du_0\|_\infty |h| = v_+(x, 0),$$

which implies the desired Lipschitz bound. \square

Remark A.6. Let $E \subset \mathbb{R}^2$ be a closed set in \mathbb{R}^2 and let $u_E(x)$ be a bounded Lipschitz continuous function such that

$$(A.4) \quad \partial E = \{x \in \mathbb{R}^2 \text{ s.t. } u_E(x) = 0\} = \partial\{x \in \mathbb{R}^2 \text{ s.t. } u_E(x) > 0\} \quad \text{and} \quad E = \{x \in \mathbb{R}^2 \text{ s.t. } u_E(x) \geq 0\}.$$

Let u_E be the unique viscosity solution to (A.3) with initial datum u_E and define

$$E^+(t) := \{x \in \mathbb{R}^2 \text{ s.t. } u_E(x, t) \geq 0\} \quad \text{and} \quad E^-(t) := \{x \in \mathbb{R}^2 \text{ s.t. } u_E(x, t) > 0\}.$$

The level set flow is defined as

$$\Sigma_E(t) = \{x \in \mathbb{R}^2 \text{ s.t. } |u_E(x, t) = 0\}.$$

Due to the fact that the operator in (A.3) is geometric, which means that if u is a subsolution (resp. a supersolution) then also $f(u)$ is a subsolution (resp. a supersolution) for all monotone increasing functions f , the following result holds: if v_0 is a Lipschitz continuous function which satisfies (A.4) and v is the viscosity solution to (A.3) with initial datum v_0 , then $E^+(t) = \{x \in \mathbb{R}^2 \text{ s.t. } v(x, t) \geq 0\}$ and $E^-(t) = \{x \in \mathbb{R}^2 \text{ s.t. } v(x, t) > 0\}$.

In particular, the inner flow, the outer flow and the level set flow do not depend on the choice of the initial datum u_E but only on the set E .

Remark A.7. In the setting of Remark A.2, one can show that $u(x, t) = R(t) - |x|$, for $t \in [0, T_R)$, is a viscosity solution to (A.3). Therefore, in this case we have that $E = B_R$ and $E^+(t) = \overline{E^-(t)} = B_{R(t)}$.

An important consequence of the Comparison Principle stated in Theorem A.5, is the following result (in which we also use the distance notations introduced in (1.35) and (1.34)).

Corollary A.8.

- i) *Let $F \subset E$ two closed sets in \mathbb{R}^2 such that $d(F, E) = \delta > 0$. Then $F^+(t) \subset E^-(t)$ for all $t > 0$, and the map $t \rightarrow d(F^+(t), E^-(t))$ is nondecreasing.*
- ii) *Let $v : \mathbb{R}^2 \times [0, T) \rightarrow \mathbb{R}$ be a bounded uniformly continuous viscosity supersolution to (A.3), and assume that*

$$F \subseteq \{x \in \mathbb{R}^2 \text{ s.t. } v(x, 0) \geq 0\}.$$

Then

$$F^+(t) \subseteq \{x \in \mathbb{R}^2 \text{ s.t. } v(x, t) \geq 0\},$$

for all $t \in (0, T)$.

Moreover, if

$$d(F, \{x \in \mathbb{R}^2 \text{ s.t. } v(x, 0) > 0\}) = \delta > 0,$$

then

$$F^+(t) \subseteq \{x \in \mathbb{R}^2 \text{ s.t. } v(x, t) > 0\},$$

for all $t \in (0, T)$, and

$$d\left(F^+(t), \{x \in \mathbb{R}^2 \text{ s.t. } v(x, t) > 0\}\right) \geq \delta.$$

iii) Let $w : \mathbb{R}^2 \times [0, T) \rightarrow \mathbb{R}$ be a bounded uniformly continuous viscosity subsolution to (A.3), and assume that

$$E \supseteq \{x \in \mathbb{R}^2 \text{ s.t. } w(x, 0) \geq 0\}.$$

Then

$$E^+(t) \supseteq \{x \in \mathbb{R}^2 \text{ s.t. } w(x, t) \geq 0\},$$

for all $t \in (0, T)$.

Moreover, if

$$d(E, \{x \in \mathbb{R}^2 \text{ s.t. } w(x, 0) \geq 0\}) = \delta > 0,$$

then

$$E^-(t) \supseteq \{x \in \mathbb{R}^2 \text{ s.t. } w(x, t) \geq 0\},$$

for all $t \in (0, T)$, and

$$d\left(E^-(t), \{x \in \mathbb{R}^2 \text{ s.t. } w(x, t) \geq 0\}\right) \geq \delta.$$

Proof. First, we prove i). Since $F \subseteq E$ and $d(E, F) = \delta$, then it is easy to check that $d_E(x) \geq d_F(x) + \delta$.

Let now $C > 2\delta$ and define

$$u_F(x) := \max\{-C - \delta, \min\{d_F(x), C - \delta\}\} \quad \text{and} \quad u_E(x) := \max\{-C, \min\{d_E(x), C\}\}.$$

So, again we obtain that $u_F(x) + \delta \leq u_E(x)$. Therefore, by the Comparison Principle in Theorem A.5, we get that $u_E(x, t) \geq u_F(x, t) + \delta$ for every $t > 0$.

This in turn implies that $F^+(t) \subseteq E^-(t)$ and moreover that $d(F^+(t), E^-(t)) \geq \delta$, due to the fact that $u_E(x, t)$ and $u_F(x, t)$ are 1-Lipschitz in x , by Theorem A.5.

If we repeat the same argument with initial data $E^-(t)$ and $F^+(t)$, we obtain the desired statement in i).

We prove now ii). For this, we distinguish two cases: if

$$(A.5) \quad d(F, \{x \in \mathbb{R}^2 \text{ s.t. } v(x, 0) > 0\}) = \delta > 0,$$

we let

$$E := \overline{\{x \in \mathbb{R}^2 \text{ s.t. } v(x, 0) > 0\}}.$$

Then, by item i), we get that $F^+(t) \subseteq E^-(t)$ and $d(F^+(t), E^-(t)) \geq \delta$. Let u_E be the unique viscosity solution to (A.3) with $u_E(x, 0) = v(x, 0)$. Then, by the Comparison Principle in Theorem A.5, we get that $u_E(x, t) \leq v(x, t)$ for all $t \in (0, T)$. In turn, this implies that $E^-(t) \subseteq \{x \in \mathbb{R}^2 \text{ s.t. } v(x, t) > 0\}$, and this permits to conclude that ii) holds true, under the assumption in (A.5).

If, on the other hand, we have that (A.5) does not hold, we write

$$d(F, \{x \in \mathbb{R}^2 \text{ s.t. } v(x, 0) > 0\}) \geq 0.$$

Then, by the uniform continuity of $v(\cdot, 0)$, we have that for every $\varepsilon > 0$ there exists $\delta_\varepsilon > 0$ such that

$$d(F, \{x \in \mathbb{R}^2 \text{ s.t. } v(x, 0) > -\varepsilon\}) \geq \delta_\varepsilon > 0.$$

So we repeat the argument above (based on (A.5)) substituting $v(x, t)$ with the function $v(x, t) + \varepsilon$ and E with $\overline{\{x \in \mathbb{R}^2 \text{ s.t. } v(x, 0) > -\varepsilon\}}$. This gives that $F^+(t) \subseteq E^-(t)$, and $E^-(t) \subseteq \{x \in \mathbb{R}^2 \text{ s.t. } v(x, t) > -\varepsilon\}$ for all $\varepsilon > 0$. Therefore $F^+(t) \subseteq \{x \in \mathbb{R}^2 \text{ s.t. } v(x, t) \geq 0\}$.

This completes the proof of ii). The proof of iii) is completely analogous, and we omit it. \square

Remark A.9. Observe that if E is a compact set and in particular $E \subseteq B_R$ for some $R > 0$, then by Remark A.7 and Corollary A.8 we have that $E^+(t) \subseteq B_{R(t)}$ where $R(t) < R$ has been defined in Remark A.2. In particular, there exists $T_E \leq T_R$ such that $T_E = \sup\{t > 0 \text{ s.t. } \text{int } E^+(t) \neq \emptyset\}$.

Now, we define the lower and upper semicontinuous envelopes of a family of sets $C(t) \subseteq \mathbb{R}^2$ as follows:

$$C_\star(t) := \bigcup_{\varepsilon > 0} \bigcap_{0 \leq t - \varepsilon < s < t + \varepsilon} C(s) \quad \text{and} \quad C^\star(t) := \bigcap_{\varepsilon > 0} \bigcup_{0 \leq t - \varepsilon < s < t + \varepsilon} C(s).$$

We have that $C_\star(t) \subseteq C(t) \subseteq C^\star(t)$. Moreover for any sequence $(x_n, t_n) \rightarrow (x, t)$, if $x_n \in \overline{C^\star(t_n)}$ then $x \in \overline{C^\star(t)}$, whereas, if $x_n \notin \text{int}(C_\star(t_n))$, then $x \notin \text{int}(C_\star(t))$.

If $C^\star(t) = C(t) = C_\star(t)$ for every t , we say that the family is continuous.

We also need a result to compare geometric sub and supersolutions to (1.1) with the level set flow, see [8].

Proposition A.10. *Let $C(t) \subseteq \mathbb{R}^2$ for $t \in [0, T]$, be a continuous family of sets with compact Lipschitz boundaries, which are piecewise of class $C^{1,1}$ outside a finite number of angular² points.*

Fix $E \subset \mathbb{R}^2$ and u_E a bounded Lipschitz continuous function such that

$$E = \{x \in \mathbb{R}^2 \text{ s.t. } u_E(x) \geq 0\} \quad \text{and} \quad \partial E = \{x \in \mathbb{R}^2 \text{ s.t. } u_E(x) = 0\}.$$

Consider the inner and outer flows associated to E , according to (1.8).

i) *Assume that there exists $\delta > 0$ such that at every $x \in \partial C(t)$ where $\partial C(t)$ is $C^{1,1}$ there holds*

$$(A.6) \quad \partial_t x \cdot \nu(x) \geq -H_{C(t)}^K(x) + \delta.$$

Moreover, assume that

$$(A.7) \quad \begin{aligned} &\text{at every angular point } x \in \partial C(t) \text{ there exists } r_0 > 0 \text{ such that} \\ &\text{the set } B(x, r) \cap C(t) \text{ is convex for all } r < r_0. \end{aligned}$$

Then, if $E \subseteq C(0)$, with $d(E, C(0)) = k \geq 0$, it holds that $E^+(t) \subset C(t)$ for all $t \in [0, T]$, with $d(E^+(t), C(t)) \geq k \geq 0$.

ii) *Assume that there exists $\delta > 0$ such that at every $x \in \partial C(t)$ where $\partial C(t)$ is $C^{1,1}$ it holds*

$$(A.8) \quad \partial_t x \cdot \nu(x) \leq -H_{C(t)}^K(x) - \delta.$$

Moreover, assume that

$$(A.9) \quad \begin{aligned} &\text{at every point angular } x \in \partial C(t) \text{ there exists } r_0 > 0 \text{ such that} \\ &\text{the set } B(x, r) \cap (\mathbb{R}^2 \setminus C(t)) \text{ is convex for all } r < r_0. \end{aligned}$$

Then, if $E \supseteq C(0)$, it holds that $E^-(t) \supseteq C(t)$ for all $t \in [0, T]$.

Moreover, if $d(C(0), \{x \in \mathbb{R}^2 \text{ s.t. } u_E(x) > 0\}) = k > 0$, it holds that $E^-(t) \supset C(t)$ for all $t \in [0, T]$, with $d(E^-(t), C(t)) \geq k$.

Proof. We give just a sketch of the proof of i), since it relies on classical arguments in viscosity solution theory and level set methods (the proof of ii) is analogous), see [8].

For $\varepsilon > 0$ sufficiently small, we define the function

$$u_\varepsilon(x, t) := \max \{0, \min\{\varepsilon, d_{C(t)}(x)\}\}.$$

We claim that for $\varepsilon > 0$ sufficiently small (depending on δ in (A.6)) the function u_ε is a viscosity supersolution to (A.3). If the claim is true, then the statement in i) is a direct consequence of the Comparison Principle in Corollary A.8.

To prove the claim, for every $\lambda \in [0, \varepsilon]$, we define

$$C_\lambda(t) := \{x \in C(t) \text{ s.t. } d_{C(t)}(x) \geq \lambda\}.$$

Note that $u_\varepsilon = 0$ on $\overline{\mathbb{R}^2 \setminus C(t)}$, $u_\varepsilon = \lambda$ on $\partial C_\lambda(t)$ and $u_\varepsilon = \varepsilon$ on $C_\varepsilon(t)$.

Due to the regularity assumption on $C(t)$, we have that for every $\lambda \in [0, \varepsilon]$, the sets $C_\lambda(t)$ are Lipschitz continuous, piecewise $C^{1,1}$ outside a finite number of angular points and satisfy the following property: at every angular point $x \in \partial C_\lambda(t)$ there exists $r_0 > 0$ such that the set $B(x, r) \cap C_\lambda(t)$ is convex for all $r < r_0$. Therefore assumption (A.7) is satisfied for every C_λ , with $\lambda \in [0, \varepsilon]$.

Let now $x_\varepsilon \in \partial C_\varepsilon(t)$ where $C_\varepsilon(t)$ is locally of class $C^{1,1}$. Let $x_0 \in \partial C(t)$ be the unique point such that $|x_0 - x_\varepsilon| = \varepsilon$ and define

$$(A.10) \quad x_\lambda = x_0 + \frac{\lambda}{\varepsilon}(x_\varepsilon - x_0) \quad \lambda \in [0, \varepsilon].$$

First of all we observe that $H_{C_\lambda(t)}^K(x_\lambda)$ is well defined, for every $\lambda \in [0, \varepsilon]$: indeed due to assumption (A.7), we get that the sets $C_\lambda(t)$ are locally of class $C^{1,1}$ at x_λ uniformly in λ . Then we use the continuity of the K -curvature as $\lambda \rightarrow 0$ (see [8]) to see that there exists $\eta = \eta(\varepsilon) > 0$ such that

$$(A.11) \quad |H_{C_\lambda(t)}^K(x_\lambda) - H_{C(t)}^K(x_0)| \leq \eta \quad \text{for all } \lambda \in [0, \varepsilon].$$

²As customary, a point of a piecewise $C^{1,1}$ curve is called ‘‘angular’’ if the tangent directions from the two sides are different.

Finally we point out that η should depend on ε and x_0 , but actually by compactness of

$$\bigcup_{t \in [0, T]} \partial C(t),$$

we can choose it independent of x_0 .

Now we distinguish different cases according to the position of the point x , in order to prove that u_ε is a viscosity supersolution to (A.3).

If $x \in \text{int}(\mathbb{R}^2 \setminus C(t))$, or $x \in \text{int}(C_\varepsilon(t))$, then actually the equation in (A.3) is trivially satisfied since $Du_\varepsilon(x, t) = 0$ and $\partial_t u_\varepsilon(x, t) = 0$ by the continuity properties of the families $C(t)$ and $C_\varepsilon(t)$.

Now we suppose that $x \in \partial C_\varepsilon(t)$. Then it is easy to show that the set of test functions is empty, so again the equation in (A.3) is trivially satisfied.

We finally assume that $x \in \partial C_\lambda(t)$ for some $\lambda \in [0, \varepsilon]$. Observe that at every angular point $x \in \partial C_\lambda(t)$, by the assumption (A.7) (which holds also for $C_\lambda(t)$ as proved above), the set of test functions is empty so the equation in (A.3) is trivially satisfied. So assume that $C_\lambda(t)$ is locally of class $C^{1,1}$ around x . We fix $x_0 \in \partial C(t)$ such that $|x - x_0| = \lambda$. So, if $\nu(x)$ is the outer normal to $\partial C_\lambda(t)$ at x , then $\nu(x) = \frac{x_0 - x}{|x - x_0|}$ and $\nu(x) = \nu(x_0)$, so it coincides with the outer normal to $\partial C(t)$ at x_0 and $\partial_t x_0 \cdot \nu(x_0) = \partial_t x \cdot \nu(x)$. Moreover, due to (A.11), there exists $\eta = \eta(\varepsilon) > 0$ such that

$$(A.12) \quad H_{C_\lambda(t)}^K(x) \geq H_{C(t)}^K(x_0) - \eta.$$

Let ϕ be a test function for u_ε at (x, t) , then $D\phi(x, t) = -\rho\nu(x)$ for some $\rho \in [0, 1]$ for $\lambda = 0$ and $D\phi(x, t) = -\nu(x)$ for $\lambda > 0$, whereas $\phi_t(x, t) = \rho\partial_t x \cdot \nu(x)$ (with $\rho = 1$ as $\lambda > 0$). Moreover

$$(A.13) \quad H_{C_\lambda(t)}^K(x) = H_{\{y | u_\varepsilon(y, t) \geq \lambda\}}^K(x) \leq H_{\{y | \phi(y, t) > \lambda\}}^K(x).$$

Therefore, computing the equation at (x, t) , we get, using (A.12), (A.13) and (A.6),

$$\partial_t \phi(x, t) + |D\phi(x, t)| H_{\{y | \phi(y, t) > \phi(x, t)\}}^K(x) \geq \rho\partial_t x \cdot \nu(x) + \rho H_{C_\lambda(t)}^K(x) \geq \rho\partial_t x_0 \cdot \nu(x_0) + \rho H_{C(t)}^K(x_0) - \rho\eta \geq \rho(\delta - \eta).$$

So, if we choose $\varepsilon > 0$ sufficiently small, according to δ , so that $\eta(\varepsilon) \leq \delta$, then the previous inequality gives that u_ε is a supersolution to (A.3), as we claimed. \square

Now we present the following extension to the noncompact case of Proposition A.10.

Corollary A.11. *Let $C(t) \subseteq \mathbb{R}^2$ for $t \in [0, T]$, be a continuous family of sets with Lipschitz boundaries, which are piecewise of class $C^{1,1}$ outside a finite number of angular points, and such that there exists $R > 0$ such that $C(t) \cap (\mathbb{R}^2 \setminus B_R)$ is of class $C^{1,1}$ for all t .*

Fix $E \subset \mathbb{R}^2$ and u_E a bounded Lipschitz continuous function such that

$$E = \{x \in \mathbb{R}^2 \text{ s.t. } u_E(x) \geq 0\} \quad \text{and} \quad \partial E = \{x \in \mathbb{R}^2 \text{ s.t. } u_E(x) = 0\},$$

and consider the inner and outer flows associated to E , according to (1.8).

- i) *Assume that there exists $\delta > 0$ such that (A.6) holds for every $x \in \partial C(t) \cap B_{4R}$. Suppose also that (A.7) holds true.*

Moreover, assume that there exists λ_0 such that, for all $\lambda \in [0, \lambda_0]$, it holds that

$$(A.14) \quad \partial_t x \cdot \nu(x) \geq -H_{C_\lambda(t)}^K(x)$$

for all $x \in \partial C_\lambda(t) \cap (\mathbb{R}^2 \setminus B_{2R})$, where

$$C_\lambda(t) := \{x \in C(t) \text{ s.t. } d_{C(t)}(x) \geq \lambda\}.$$

Then, if $E \subset C(0)$, with $d(E, C(0)) = k \geq 0$, it holds that $E^+(t) \subset C(t)$ for all $t > 0$, with $d(E^+(t), C(t)) \geq k \geq 0$.

- ii) *Assume that there exists $\delta > 0$ such that (A.8) holds for every $x \in \partial C(t) \cap B_{4R}$. Suppose also that (A.9) holds true.*

Moreover, assume that there exists λ_0 such that for all $\lambda \in [0, \lambda_0]$, it holds that

$$(A.15) \quad \partial_t x \cdot \nu(x) \leq -H_{C_\lambda(t)}^K(x)$$

for all $x \in \partial C^\lambda(t) \cap (\mathbb{R}^2 \setminus B_{2R})$ where

$$C^\lambda(t) := \{x \in \mathbb{R}^2 \text{ s.t. } d_{C(t)} \geq -\lambda\}.$$

Then, if $E \supseteq C(0)$, it holds that $E^+(t) \supseteq C(t)$ for all $t > 0$.

In addition, if $d(C(0), \{x \in \mathbb{R}^2 \text{ s.t. } u_E(x) > 0\}) = k > 0$, it holds that $E^-(t) \supset C(t)$ for all $t > 0$, with $d(E^-(t), C(t)) \geq k$.

The proof of Corollary A.11 is similar to that of Proposition A.10, and we omit the details.

We also have the following semicontinuity type result for the outer evolutions.

Proposition A.12. *There holds*

$$(A.16) \quad \liminf_{\eta \searrow 0} |E^+(t + \eta)| \geq |\text{int } E^+(t)|.$$

Proof. We claim that for any fixed $t > 0$ and a.e. in \mathbb{R}^2 ,

$$(A.17) \quad \liminf_{\eta \searrow 0} \chi_{\{u_E(\cdot, t + \eta) \geq 0\}} \geq \chi_{\{\text{int}(\{u_E(\cdot, t) \geq 0\})\}}.$$

To show (A.17), it is enough to consider a point $x \in \text{int}(\{u_E(\cdot, t) \geq 0\})$, so that $\{u_E(\cdot, t) \geq 0\} \supset B_r(x)$ for some $r > 0$. Then, recalling formula (A.2) in Remark A.2, we have that $C(r(\eta)) = C(r) - \eta$, for $\eta \in (0, T_r)$, where $T_r > 0$ is the extinction time of the ball B_r under the flow (1.1).

Hence, by Remark A.7 and Corollary A.8 we get

$$\{u_E(\cdot, t + \eta) \geq 0\} \supset B_{r(\eta)}(x), \quad \text{for all } \eta \in (0, T_r).$$

In particular, it follows that

$$\liminf_{\eta \searrow 0} u_E(x, t + \eta) \geq 0, \quad \text{for all } x \in \text{int}\{u_E(\cdot, t) \geq 0\},$$

which implies (A.17).

Then, by (A.17) and the Fatou Lemma, for all $t > 0$ we obtain

$$\begin{aligned} \liminf_{\eta \searrow 0} |E^+(t + \eta)| &= \liminf_{\eta \searrow 0} \int_{\mathbb{R}^2} \chi_{\{u_E(\cdot, t + \eta) \geq 0\}}(x) dx \\ &\geq \int_{\mathbb{R}^2} \liminf_{\eta \searrow 0} \chi_{\{u_E(\cdot, t + \eta) \geq 0\}}(x) dx \geq |\text{int}(\{u_E(\cdot, t) \geq 0\})| = |\text{int } E^+(t)|, \end{aligned}$$

establishing (A.16). \square

In the case of homogeneous kernels, i.e. under the assumption in (1.27), the geometric flow possesses a useful time scaling property as follows.

Lemma A.13. *Assume that $K(x) = \frac{1}{|x|^{2+s}}$ for some $s \in (0, 1)$. Let $\lambda > 0$, $M > 0$, $E \subseteq \mathbb{R}^2$ and $u_{E,\lambda}(x, t)$ be the viscosity solution to (A.3) with initial condition given by*

$$u_{E,\lambda}(x) := \max \{ -\lambda M, \min \{ d_{\lambda E}(x), \lambda M \} \}.$$

Let also $E_\lambda^+(t) := \{x \in \mathbb{R}^2 \text{ s.t. } u_{E,\lambda}(x, t) \geq 0\}$ and $E_\lambda^-(t) := \{x \in \mathbb{R}^2 \text{ s.t. } u_{E,\lambda}(x, t) > 0\}$. Then

$$E_\lambda^\pm(t) = \lambda E_1^\pm \left(\frac{t}{\lambda^{1+s}} \right).$$

Proof. For every $x \in \mathbb{R}^2$ such that $-M \leq d_E(x) \leq M$, we have that

$$\lambda u_{E,1}(x) = \lambda d_E(x) = d_{\lambda E}(\lambda x) = u_{E,\lambda}(\lambda x).$$

Moreover if $d_E(x) \geq M$, then $\lambda u_{E,1}(x) = \lambda M = u_{E,\lambda}(\lambda x)$, and analogously for $d_E(x) \leq -M$. Therefore we get that $\lambda u_{E,1}(\frac{x}{\lambda}) = u_{E,\lambda}(x)$. Moreover, by the scaling properties of K , we have that $H_E^s(x) = \lambda^{-s} H_{\lambda E}^s(\lambda x)$. Therefore the function $\lambda u_{E,1}(\frac{x}{\lambda}, \frac{t}{\lambda^{1+s}})$ is a viscosity solution to (A.3), with initial datum $u_{E,\lambda}(x)$. By the uniqueness of viscosity solutions, given in Theorem A.5, we get that $\lambda u_{E,1}(\frac{x}{\lambda}, \frac{t}{\lambda^{1+s}}) = u_{E,\lambda}(x, t)$. From this we deduce the desired statement. \square

Acknowledgements. This work has been supported by the Australian Research Council Discovery Project DP170104880 “NEW – Nonlocal Equations at Work”, and by the University of Pisa Project PRA 2017 “Problemi di ottimizzazione e di evoluzione in ambito variazionale”. The authors are members of the INdAM-GNAMPA.

REFERENCES

- [1] G. Barles, H. M. Soner, and P. E. Souganidis, *Front propagation and phase field theory*, SIAM J. Control Optim. **31** (1993), no. 2, 439–469, DOI 10.1137/0331021. MR1205984
- [2] Giovanni Bellettini, Vicent Caselles, Antonin Chambolle, and Matteo Novaga, *Crystalline mean curvature flow of convex sets*, Arch. Ration. Mech. Anal. **179** (2006), no. 1, 109–152, DOI 10.1007/s00205-005-0387-0. MR2208291
- [3] Giovanni Bellettini and Maurizio Paolini, *Two examples of fattening for the curvature flow with a driving force*, Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl. **5** (1994), no. 3, 229–236. MR1298266
- [4] Samuel Biton, Pierre Cardaliaguet, and Olivier Ley, *Nonfattening condition for the generalized evolution by mean curvature and applications*, Interfaces Free Bound. **10** (2008), no. 1, 1–14, DOI 10.4171/IFB/177. MR2383534
- [5] Claudia Bucur and Enrico Valdinoci, *Nonlocal diffusion and applications*, Lecture Notes of the Unione Matematica Italiana, vol. 20, Springer; Unione Matematica Italiana, Bologna, 2016. MR3469920
- [6] Luis Caffarelli, Jean-Michel Roquejoffre, and Ovidiu Savin, *Nonlocal minimal surfaces*, Comm. Pure Appl. Math. **63** (2010), no. 9, 1111–1144, DOI 10.1002/cpa.20331. MR2675483
- [7] Annalisa Cesaroni and Matteo Novaga, *The isoperimetric problem for nonlocal perimeters*, Discrete Contin. Dyn. Syst. Ser. S **11** (2018), no. 3, 425–440, DOI 10.3934/dcds.2018023. MR3732175
- [8] Antonin Chambolle, Massimiliano Morini, and Marcello Ponsiglione, *Nonlocal curvature flows*, Arch. Ration. Mech. Anal. **218** (2015), no. 3, 1263–1329, DOI 10.1007/s00205-015-0880-z. MR3401008
- [9] Antonin Chambolle, Matteo Novaga, and Berardo Ruffini, *Some results on anisotropic fractional mean curvature flows*, Interfaces Free Bound. **19** (2017), no. 3, 393–415, DOI 10.4171/IFB/387. MR3713894
- [10] Yun Gang Chen, Yoshikazu Giga, and Shun'ichi Goto, *Uniqueness and existence of viscosity solutions of generalized mean curvature flow equations*, J. Differential Geom. **33** (1991), no. 3, 749–786. MR1100211
- [11] Eleonora Cinti, Joaquim Serra, and Enrico Valdinoci, *Quantitative flatness results and BV-estimates for stable nonlocal minimal surfaces*, J. Differential Geom. (2018).
- [12] Eleonora Cinti, Carlo Sinestrari, and Enrico Valdinoci, *Neckpinch singularities in fractional mean curvature flows*, Proc. Amer. Math. Soc. **146** (2018), no. 6, 2637–2646, DOI 10.1090/proc/14002. MR3778164
- [13] L. C. Evans and J. Spruck, *Motion of level sets by mean curvature. I*, J. Differential Geom. **33** (1991), no. 3, 635–681. MR1100206
- [14] Yoshikazu Giga, *Surface evolution equations. A level set approach*, Monographs in Mathematics, vol. 99, Birkhäuser Verlag, Basel, 2006. MR2238463
- [15] Tom Ilmanen, *Generalized flow of sets by mean curvature on a manifold*, Indiana Univ. Math. J. **41** (1992), no. 3, 671–705, DOI 10.1512/iumj.1992.41.41036. MR1189906
- [16] Cyril Imbert, *Level set approach for fractional mean curvature flows*, Interfaces Free Bound. **11** (2009), no. 1, 153–176, DOI 10.4171/IFB/207. MR2487027
- [17] Vladimir Maz'ya, *Lectures on isoperimetric and isocapacitary inequalities in the theory of Sobolev spaces*, Heat kernels and analysis on manifolds, graphs, and metric spaces (Paris, 2002), Contemp. Math., vol. 338, Amer. Math. Soc., Providence, RI, 2003, pp. 307–340, DOI 10.1090/conm/338/06078. MR2039959
- [18] Mariel Sáez and Enrico Valdinoci, *On the evolution by fractional mean curvature*, Comm. Anal. Geom. **27** (2019), no. 1.
- [19] Ovidiu Savin and Enrico Valdinoci, *Regularity of nonlocal minimal cones in dimension 2*, Calc. Var. Partial Differential Equations **48** (2013), no. 1-2, 33–39, DOI 10.1007/s00526-012-0539-7. MR3090533
- [20] Halil Mete Soner, *Motion of a set by the curvature of its boundary*, J. Differential Equations **101** (1993), no. 2, 313–372, DOI 10.1006/jdeq.1993.1015. MR1204331